

Chapter 4

Field Quantization

General idea of field quantization: promote field amplitudes to operators on suitable Hilbert space. For the Schrödinger field: 'second quantization'. For the Maxwell field: 'first quantization'.

Hierarchy of theories		
Classical mechanics point particles classical path Geometrical optics	Wave mechanics wave function matter waves wave optics	Quantum field theory wave/field operator particle creation/annihilation photons

Table 4.1:

Reminder: harmonic oscillator, two degrees of freedom

$$H = \frac{p_a^2}{2m} + \frac{K}{2}q_a^2 + \frac{p_b^2}{2m} + \frac{K}{2}q_b^2 \quad (4.1)$$

eigenfrequencies $\omega_a = \omega_b = \sqrt{K/m}$. Quantization in terms of creation and annihilation operators:

$$q_a = \sqrt{\frac{\hbar}{2m\omega_a}}(a + a^\dagger), \quad p_a = -i\sqrt{\frac{\hbar m\omega_a}{2}}(a - a^\dagger) \quad (4.2)$$

$$a = \sqrt{\frac{m\omega_a}{2\hbar}}\left(q_a + \frac{i}{m\omega_a}p_a\right), \quad a^\dagger = \sqrt{\frac{m\omega_a}{2\hbar}}\left(q_a - \frac{i}{m\omega_a}p_a\right) \quad (4.3)$$

and b, b^\dagger for q_b, p_b . Commutation relations

$$[a, a^\dagger] = 1, \quad [b, b^\dagger] = 1 \quad (4.4)$$

all others are zero.

'Number operator' $a^\dagger a$ counts excitations. In quantum field theory, an excitation is interpreted as a particle, and $a^\dagger a$ counts particles with quantum numbers (mode function, spin etc.) specific to the oscillator of type 'a'. In other words: a 'particle' is an excitation of a 'normal mode' of a field that behaves like a harmonic oscillator. The operators a^\dagger and a implement the creation/annihilation of this particle.

4.1 Basics: phonons

This is a simple model from classical mechanics to start with. We avoid technical complications and work with a system made up of discrete oscillators and of finite size. The basic observation that we are going to make: the 'small amplitude vibrations' of classical mechanics become the 'modes' of a 'field' whose dispersion relation is similar to that of a relativistic particle.

4.1.1 Hamiltonian

Hamiltonian:

$$H = \sum_n \frac{p(x_n)^2}{2m} + \sum_n \frac{K}{2} [q(x_n + a) - q(x_n)]^2 + \sum_n \frac{m\Omega^2}{2} q(x_n)^2 \quad (4.5)$$

Here, the oscillator displacement is $q(x_n)$, its equilibrium position is x_n . The conjugate momentum is $p(x_n)$. The first term in (4.5) is the kinetic energy, the second the potential energy from the restoring force to equilibrium (spring constant $m\Omega^2$), and the third the potential energy from linear coupling to the neighboring oscillator, with spring constant K . The oscillators are separated by the discrete lattice vector a – this is a one-dimensional chain. The sum runs over N oscillators in total. We take periodic boundary conditions: $q(x_{n+N}) = q(x_n + Na) = q(x_n)$.

In classical mechanics, $q_n \equiv q(x_n)$ and $p_{n'}$ are canonically conjugate. This means that the Poisson brackets are $\{q_n, p_{n'}\} = \delta_{nn'}$.

We promote this to a quantized theory by a recipe called 'canonical quantization': replace Poisson brackets by commutators. In other words: we promote coordinates and conjugate momenta to operators that do not commute:

$$[q_n, p_{n'}] = i\hbar\delta_{nn'}, \quad [q_n, q_{n'}] = 0, \quad (4.6)$$

of course, the momenta commute also with themselves.

The time evolution of observables follows in the quantized theory from the Heisenberg equations, for example:

$$\frac{dq_n}{dt} = \frac{i}{\hbar} [H, q_n] \quad (4.7)$$

with Hamilton operator H obtained from (4.5) written with operators. In classical mechanics, we have the same equation written in terms of Poisson brackets.

4.1.2 Mode expansion

The system is periodic, therefore it is useful to make a Fourier expansion:

$$q(x) = \sum_{k \in \text{BZ}} \tilde{q}(k)e^{ikx}, \quad \tilde{q}(k) = \frac{1}{N} \sum_n q(x_n)e^{-ikx_n} \quad (4.8)$$

and similarly for $p(x_n)$. The discrete wave vectors k run from $-\pi/a$ to π/a in steps of $2\pi/L$ where $L = Na$ is the length of the chain. These points make

up the 'Brillouin zone' BZ. Since the operator q is hermitean, one must have $\tilde{q}(-k) = \tilde{q}^\dagger(k)$. The commutation relations become

$$[\tilde{q}(k), \tilde{p}^\dagger(k')] = \frac{i\hbar}{N^2} \sum_n e^{i(k-k')x_n} \quad (4.9)$$

Now for discretized wave vectors in the first Brillouin zone, $-\pi/a < k \leq \pi/a$, this sum of exponentials is nonzero only if $k - k' = 0 \pmod{2\pi/a}$. In addition, the difference of k -vectors is limited to $|k - k'| \leq 2\pi(N-1)/N$. Hence only for $k = k'$, the sum gives a nonzero result (which is trivially equal to N). So we have

$$[\tilde{q}(k), \tilde{p}^\dagger(k')] = \frac{i\hbar}{N} \delta_{kk'} \quad (4.10)$$

with a Kronecker for the discrete wavenumbers.

A similar calculation gives for the Hamiltonian (see Exercise)

$$H = N \sum_{k \in \text{BZ}} \frac{1}{2m} \tilde{p}^\dagger(k) \tilde{p}(k) + N \sum_k \left(\frac{m\Omega^2}{2} + K(1 - \cos ka) \right) \tilde{q}^\dagger(k) \tilde{q}(k) \quad (4.11)$$

The key observation is that this is a sum of independent oscillators: one for each value of k . In classical terms: we have found the 'normal modes' of the system. But these modes provide also one basic recipe for field quantization: find the 'modes' that bring the Hamiltonian into a diagonal form¹ and promote the amplitudes of these modes to operators.

We note that the frequency for a mode with wavenumber k is given by

$$\omega_k^2 = \Omega^2 + \frac{2K}{m} (1 - \cos ka) \approx \Omega^2 + c^2 k^2 \quad (4.12)$$

In the last line, we have introduced the large-wavelength limit $ka \ll 1$ and expanded the cosine. This looks like the dispersion relation for a particle with rest mass Ω , with the speed of light being replaced by the 'speed of sound' $c^2 = K a^2 / m$. This is indeed the speed of long-wavelength oscillations that propagate on the chain. In the limit $\Omega \rightarrow 0$, one gets the linear dispersion $\omega_k = c|k|$ similar to light waves or sound waves.

4.1.3 Quantization

For each k , we have a harmonic oscillator and quantize in terms of the creation and annihilation operators

$$a_k = \sqrt{\frac{Nm\omega_k}{2\hbar}} \left[\tilde{q}(k) + \frac{i}{m\omega_k} \tilde{p}(k) \right], \quad a_k^\dagger = \sqrt{\frac{Nm\omega_k}{2\hbar}} \left[\tilde{q}^\dagger(k) - \frac{i}{m\omega_k} \tilde{p}^\dagger(k) \right]. \quad (4.13)$$

One can check that the commutation relations are indeed those for independent oscillators

$$[a_k, a_{k'}^\dagger] = \delta_{kk'} \quad (4.14)$$

Hamiltonian (4.11) becomes

$$H = \sum_{k \in \text{BZ}} \frac{\hbar\omega_k}{2} (a_k^\dagger a_k + a_k a_k^\dagger) = \sum_k \hbar\omega_k \left(a_k^\dagger a_k + \frac{1}{2} \right) \quad (4.15)$$

¹This could have been done at the classical level, of course.

which is the usual form for the energy of a collection of independent oscillators. This kind of result one always gets when a field is quantized. The real benefit of the quantization procedure is the following 'mode expansion' of the particle displacement that is now operator-valued:

$$q(x, t) = \sum_{k \in \text{BZ}} \sqrt{\frac{\hbar}{2m\omega_k N}} \left(a_k(t) e^{ikx} + a_k^\dagger(t) e^{-ikx} \right) \quad (4.16)$$

The t -dependence of the annihilation and creation operators is simple: multiply with $e^{\pm i\omega_k t}$.

We now discuss some applications. The physics behind this result is related to 'phonons'. These quasiparticles correspond to the excitations of the normal modes we just found. A typical stationary state of the system is specified by a set of occupation numbers $\{n_k\}$, one for each plane-wave mode $k \in \text{BZ}$. In this state, one says that a number n_k of phonons make up the excitation of the mode k . There is no contradiction in having many phonons exciting a given mode: phonons are 'bosons' that are not subject to Pauli's exclusion principle.

4.1.4 Applications

Zero-point oscillations

The ground state of the quantized oscillator chain is characterized by the equations $a_k |vac\rangle = 0$. Equivalently, $n_k = 0$ for all normal modes k . The mean value of the displacement in the ground state is zero:

$$\langle q(x) \rangle_{vac} = \sum_{k \in \text{BZ}} (\dots) \langle a_k \rangle_{vac} \quad (4.17)$$

because $\langle a_k \rangle_{vac} = 0$ in the ground state (in any stationary state, in fact).

But the variance of $q(x)$ is not zero: this is related to the zero-point fluctuations of each phonon mode being a quantized harmonic oscillator. These zero-point oscillations can be detected with scattering experiments. We single out artificially one oscillator from the chain and compute the amplitude for a scattered wave with a wave vector transfer p :

$$\langle e^{ipq_n} \rangle = e^{-\frac{1}{2} p^2 \langle q_n^2 \rangle}$$

This result is correct for a gaussian statistics² of the variable q_n . This average amplitude can be measured by letting the scattered wave interfere with a reference wave. The diffraction of X-rays by a crystal also leads to a similar behaviour of the intensities diffracted into a given order. In that case, p is the wave vector transfer of the diffraction peak.

The computation of the variance gives

$$\begin{aligned} \langle q_n^2 \rangle_{vac} &= \sum_k \frac{\hbar}{2m\omega_k N} \langle a_k e^{ikx_n} a_k^\dagger e^{-ikx_n} \rangle_{vac} \\ &= a \int_{\text{BZ}} \frac{dk}{2\pi} \frac{\hbar}{2m\omega_k}, \end{aligned} \quad (4.18)$$

²We are a cheating a little bit, since q_n is in fact an operator, but one can introduce a 'quasi-distribution' that is indeed gaussian in the ground state (and at finite temperature as well).

where we used the standard rule of replacing a sum over k into an integral for a system with large length $L = Na$. This integral has some value that depends on $\Omega a/c$:

$$\langle q_n^2 \rangle_{\text{vac}} = \frac{\hbar a}{2mc} \int_{-\pi}^{\pi} \frac{dx}{2\pi} [(\Omega a/c)^2 + 2(1 - \cos x)]^{-1/2} \quad (4.19)$$

The integral is plotted in Figure 4.1. The long-wavelength approximation, $\omega_k^2 =$




Figure 4.1: Two upper curves: integral in Eq.(4.19) as a function of $\Omega a/c$. The limiting value for $\Omega \rightarrow 0$ is ≈ 1.18 . The dashed line corresponds to the approximation $1 - \cos x \approx x^2/2$ under the integral. Dotted line: 3D equivalent.

$\Omega^2 + c^2 k^2$ gives the dashed line which is in good overall agreement. We shall use this approximation in the following because it simplifies the calculations.

Three-dimensional case

We generalize to three dimensions: now, the number N of oscillators scales with volume L^3 of the system. The sum over \mathbf{k} runs over a 3D Brillouin zone. The phonon dispersion relation for a simple cubic lattice is given by

$$\omega_{\mathbf{k}}^2 = \Omega^2 + \frac{2K}{m} (3 - \cos k_x a - \cos k_y a - \cos k_z a) \approx \Omega^2 + c^2 \mathbf{k}^2 \quad (4.20)$$

with the sound velocity $c^2 = Ka^2/m$ as before. In the long-wavelength limit, the dispersion relation is isotropic.

A similar calculation leads to the zero-point fluctuations

$$\langle q_n^2 \rangle_{\text{vac}} = a^3 \int_{\text{BZ}} \frac{d^3 k}{(2\pi)^3} \frac{\hbar}{2m\omega_{\mathbf{k}}} \quad (4.21)$$

Let us make an approximation that is often used, and consider a spherical Brillouin zone with the same number of points/the same volume. Its radius k_{max} is fixed by

$$\frac{4\pi}{3} k_{\text{max}}^3 = \left(\frac{2\pi}{a}\right)^3 \quad (4.22)$$

where the right hand side is the number of points in the 'Brillouin cube' of the 3D cubic lattice. We write $k_{\text{max}} = \omega_D/c$ in terms of the so-called Debye frequency. This quantity is often used when $\Omega = 0$ (linear phonon dispersion

relation). Typical values of the Debye frequency are of order 10^{12} s^{-1} , depending on the crystal lattice constant and stiffness.

Hence, with this approximation:

$$\langle q_n^2 \rangle_{\text{vac}} = \frac{\hbar a^3}{2m} 4\pi \int_0^{k_{\text{max}}} \frac{k^2 dk}{(2\pi)^3} \omega_k^{-1} \quad (4.23)$$

which gives another integral that depends on $\Omega a/c$ (left as an exercise). Making the long-wavelength approximation and taking a spherical Brillouin zone, the resulting zero-point fluctuation is plotted as a dotted line in Figure 4.1. The fluctuation is normalized to $\hbar a/(2mc)$, as in Eq.(4.19).

Specific heat

We are now going to make some thermodynamics. Recall the quantum thermodynamics for a single oscillator: the free energy is given by

$$\mathcal{F} = (1/\beta) \log[2 \sinh(\beta \hbar \omega/2)] \quad (4.24)$$

with $1/\beta = k_B T$ and the specific heat (per degree of freedom) is found from

$$C = \frac{\partial}{\partial T} \frac{\partial}{\partial \beta} (\beta \mathcal{F}) \quad (4.25)$$

(One differentiates the mean energy with respect to temperature.)

Now, the harmonic chain is a collection of oscillators with frequencies $\omega_{\mathbf{k}}$. In thermodynamic equilibrium, we assume that each normal mode is specified by a thermal distribution. All normal modes are independent, hence total free energy is the sum of free energies:

$$\mathcal{F} = \sum_{\mathbf{k}} (1/\beta) \log 2 \sinh(\beta \hbar \omega_{\mathbf{k}}/2) = \frac{a^3 N}{\beta} \int_{\text{BZ}} \frac{d^3 k}{(2\pi)^3} \log 2 \sinh(\beta \hbar \omega_{\mathbf{k}}/2) \quad (4.26)$$

This is proportional to the volume $a^3 N$ of the system, as it should be for the extensive quantity 'free energy'. We thus define a density of free energy $f = \mathcal{F}/a^3 N$.

This free energy density contains a zero-point contribution plus a thermal one. We compute first the zero-point contribution for a linear dispersion $\omega_{\mathbf{k}} = c|\mathbf{k}|$:

$$\begin{aligned} f(T \rightarrow 0) &= \frac{4\pi}{\beta} \int_0^{k_{\text{max}}} \frac{k^2 dk}{(2\pi)^3} \frac{\beta \hbar \omega_{\mathbf{k}}}{2} \\ &\approx 2\pi \hbar c \int_0^{k_{\text{max}}} \frac{k^3 dk}{(2\pi)^3} = \frac{\hbar c k_{\text{max}}^4}{(4\pi)^2} \sim \frac{\hbar c}{a^4} \end{aligned} \quad (4.27)$$

This is temperature-independent (by construction), hence it does not contribute to the specific heat. The result (4.27) corresponds to the energy of one phonon at the edge of the Brillouin zone, $\hbar c/a \sim \hbar \omega_D$, per unit cell of the

crystal. Of course, this zero-point energy is not directly observable, one would have to destroy the crystal to see it. But sometimes corrections due to finite surface size, thickness and so on appear that can be measured as ‘‘Casimir forces’’. However, the main field that contributes in that context is the electromagnetic one, not the phonon field.³

Finally, the temperature-dependent part of the free energy is the one that is relevant for the specific heat (4.25) as well. This was pointed out and calculated by Einstein and Debye, using different models for the distribution of phonon frequencies. With our simple model $\omega_k = c|k|$, we get

$$\begin{aligned} f(T) - f(0) &\approx \frac{4\pi}{\beta} \int_0^{k_{\max}} \frac{k^2 dk}{(2\pi)^3} \log(1 - e^{-\beta\hbar ck}) \\ &\approx \frac{1}{\beta^4 \hbar^3 c^3} \frac{4\pi}{(2\pi)^3} \int_0^{\beta\hbar\omega_D} x^2 dx \log(1 - e^{-x}) \end{aligned} \quad (4.28)$$

This result is plotted in Figure 4.2 in units of $(\beta^4 \hbar^3 c^3)^{-1}$ as a function of




Figure 4.2: Free energy per unit volume of a harmonic lattice in 3D, Eq.(4.28), in units of $(\beta^4 \hbar^3 c^3)^{-1}$ and as a function of $T/T_D = (\beta\hbar\omega_D)^{-1}$. The phonon dispersion relation is approximated as $\omega_k = c|k|$. The value for $T \rightarrow 0$ is $-\pi^2/90 \approx -0.11$.

the scaled temperature $(\beta\hbar\omega_D)^{-1} = k_B T / \hbar\omega_D$ (Debye temperature typically $\hbar\omega_D/k_B \sim 300$ K). At low temperatures, $k_B T \ll \hbar\omega_D$, the integral is constant. In this regime, the free energy (4.28) scales like $1/\beta^4 = T^4$. It follows, that the specific heat (4.25) per volume is proportional to T^3 in this regime. This is an important result that can be directly compared to experiments. As the temperature increases, the dependence is getting more complicated. In this ‘classical regime’, our linear chain is in fact no longer a realistic model because nonlinear contributions to the oscillations of the crystal play a role.

³More precisely, the change in electromagnetic zero-point energy per unit area for a pair of perfectly reflecting plates with distance a scales like Eq.(4.27), with c being the speed of light.

4.2 Klein-Gordon field

We now summarize a few important properties of the quantization of the Klein-Gordon field. With respect to the phonons, we are going to consider a continuous, position-dependent quantum operator $q(x)$ or $\Phi(x)$. In addition, we have to ensure that the quantization is compatible with special relativity.

4.2.1 Continuum limit

Let us first comment on the limit that the chain of oscillators merges into a continuous field (an ‘elastic string’). We shall base the discussion on the Lagrangian formulation, and our goal will be a prescription to get the Lagrangian (2.14) for the Klein-Gordon field

$$L = \int d^3x \left[\frac{1}{2} (\partial_t \Phi)^2 - \frac{c^2}{2} (\nabla \Phi)^2 - \frac{1}{2} (mc^2/\hbar)^2 \Phi^2 \right] \quad (4.29)$$

Compared to the previous formulation, this is only valid for a real-valued field $\Phi(\mathbf{x}, t)$.

The Lagrangian for the linear chain whose Hamiltonian is (4.5) reads, generalizing to three dimensions:

$$L = \sum_{\mathbf{x}} \left[\frac{M}{2} \dot{q}(\mathbf{x})^2 - \frac{M\Omega^2}{2} q(\mathbf{x})^2 - \frac{K}{2} \sum_i (q(\mathbf{x} + \mathbf{a}_i) - q(\mathbf{x}))^2 \right] \quad (4.30)$$

Here, we have written M for the oscillator mass to avoid confusion with the mass m in the Klein-Gordon field. The vectors \mathbf{a}_i correspond to the directions and distances for the coupling between neighboring oscillators. The natural replacement rule to go from the summation over \mathbf{x} to an integral is, of course:

$$\sum_{\mathbf{x}} \mapsto \int \frac{d^3x}{a_x a_y a_z} \quad (4.31)$$

We see that the mass density $\rho \equiv M/(a_x a_y a_z)$ can be factored from the Lagrangian. Note that this common factor does not change the Euler-Lagrange equations (principle of least action). We extend $q(\mathbf{x}, t)$ to a continuous function and get

$$L \mapsto \rho \int d^3x \left[\frac{1}{2} (\partial_t q)^2 - \frac{\Omega^2}{2} q^2 - \frac{K}{2M} \sum_i (q(\mathbf{x} + \mathbf{a}_i) - q(\mathbf{x}))^2 \right] \quad (4.32)$$

Finally, in the limit $a \rightarrow 0$ (at fixed ρ), we expand $q(\mathbf{x} + \mathbf{a}_i)$ in the coupling between neighbors and get

$$[q(\mathbf{x} + \mathbf{a}_i) - q(\mathbf{x})]^2 = \mathbf{a}_i^2 \left(\frac{\partial q}{\partial x_i} \right)^2 \quad (4.33)$$

where the coordinate x_i is chosen along the direction of \mathbf{a}_i . We shall take the simplest case that $|\mathbf{a}_i| = a$ in three orthogonal directions (cubic lattice) so that

$$\sum_i \mathbf{a}_i^2 \left(\frac{\partial q}{\partial x_i} \right)^2 = a^2 (\nabla q)^2 \quad (4.34)$$

With the definition for the speed of light/sound, $c^2 = Ka^2/M$, and the replacement $\Omega \mapsto mc^2/\hbar$, we thus get the Klein-Gordon Lagrangian (4.29) if we identify the Klein-Gordon field $\Phi(\mathbf{x}, t) = \sqrt{\rho}q(\mathbf{x}, t)$.

In the continuum limit $a \rightarrow 0$, the dispersion relation (4.12) also becomes identical to the relativistic one:

$$\omega_k^2 = \Omega^2 + \frac{c^2}{2a^2} \left(3 - \sum_i \cos k_i a \right) \rightarrow (mc^2/\hbar)^2 + c^2 \mathbf{k}^2. \quad (4.35)$$

We thus have formulated a ‘mechanical model’ for the propagation of a relativistic field. Of course, the modern interpretation of the field dynamics does no longer mention the underlying oscillators, be they discrete or continuous. The Klein-Gordon field ‘lives’ in empty space. This empty space behaves like a rigid medium in the sense that the field amplitude at every space point corresponds to a harmonic oscillator with frequency $\Omega = mc^2/\hbar$. This is a very large frequency already for the electron mass.

4.2.2 Quantization

We now apply the rules for the continuum limit to translate the quantized phonon field to the relativistic case. From hereon, we put $\hbar = c = 1$.

Field operator

In the discrete summations over the Brillouin zone, the spacing of the k -vectors can be first kept the same ($2\pi/L$), if we continue to work with a finite ‘quantization volume’ and periodic boundary conditions.⁴ The upper limit of the Brillouin zone, π/a , is going to infinity in the continuum limit. Hence we now have a countable, but infinite set of plane-wave modes in the expansion of the phonon displacement operator:

$$q(\mathbf{x}, t) = \sum_{\mathbf{k} \in \text{BZ}} \frac{1}{\sqrt{2M\omega_k N}} \left(a_{\mathbf{k}}(t) e^{i\mathbf{k} \cdot \mathbf{x}} + a_{\mathbf{k}}^\dagger(t) e^{-i\mathbf{k} \cdot \mathbf{x}} \right) \quad (4.36)$$

We write $MN = (M/a^3)Na^3 = \rho V$ and see that q involves the scale factor $1/\sqrt{\rho}$. For the time-dependent Klein-Gordon operator, $\Phi(x) = \sqrt{\rho}q(x)$, we thus have

$$\Phi(x) = \sum_{\mathbf{k} \in \text{BZ}} \frac{1}{\sqrt{2\omega_k V}} \left(a_{\mathbf{k}} e^{-ik_\mu x^\mu} + a_{\mathbf{k}}^\dagger e^{ik_\mu x^\mu} \right) \quad (4.37)$$

using the obvious 4-momentum $k_\mu = (\omega_k, -\mathbf{k})$ and the time evolution of the Heisenberg operators $a_{\mathbf{k}}(t)$.

Energy and momentum operators

The quantized Hamiltonian still has the same form

$$H = \sum_{\mathbf{k} \in \text{BZ}} \omega_k \left(a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \frac{1}{2} \right) \quad (4.38)$$

⁴This can be relaxed and the theory formulated in an infinite volume. The price to pay is that the $a_{\mathbf{k}}$ become operator-valued distributions that make sense only when smeared out with L^2 -integrable functions.

For each excitation in mode \mathbf{k} , the energy increases by an amount ω_k . This corresponds to the idea that particles in quantum field theory are viewed as excitations of (continuous) fields. More formally, we have the relation

$$[H, a_{\mathbf{k}}^\dagger] = \omega_k a_{\mathbf{k}}^\dagger \quad (4.39)$$

From this, we can derive that by acting with a creation operator on the vacuum state, $a_{\mathbf{k}}^\dagger|\text{vac}\rangle$, one gets a state with an energy ω_k above the vacuum state. This is compatible with the interpretation: ‘ $a_{\mathbf{k}}^\dagger$ creates a particle with momentum \mathbf{k} ’.

The vacuum is defined by the absence of any particle, $a_{\mathbf{k}}|\text{vac}\rangle = 0$. This suggests to subtract the zero-point energy in the Hamiltonian (4.38): one thus works with $H \mapsto H - \langle \text{vac} | H | \text{vac} \rangle$ or so-called ‘normally ordered’ operators (this operator ordering is expressed by the notation $:\dots:$)

$$H \mapsto \sum_{\mathbf{k} \in \text{BZ}} \frac{\omega_k}{2} : (a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + a_{\mathbf{k}} a_{\mathbf{k}}^\dagger) : = \sum_{\mathbf{k} \in \text{BZ}} \omega_k a_{\mathbf{k}}^\dagger a_{\mathbf{k}} \quad (4.40)$$

where the operators are ordered such that the annihilation operators appear to the right. By definition, the normally ordered $:H:$ gives zero when acting on the vacuum state.

Eq.(4.39) can also be used to show that the commutator between H and the field operator (4.37) generates the equations of motion:

$$i[H, \Phi] = \partial_t \Phi \quad (4.41)$$

which is of course the Heisenberg equation for the observable Φ . This can be stated in a different way: the operator H generates the translation in time of the field operator. This is a first step towards the relativistic invariance of the theory: time and space translations are part of the Lorentz-Poincaré symmetry group that every relativistic field theory should take into account.

The space translations are generated by the total momentum operator

$$\mathbf{P} = \sum_{\mathbf{k} \in \text{BZ}} \mathbf{k} a_{\mathbf{k}}^\dagger a_{\mathbf{k}}, \quad i[\mathbf{P}, \Phi] = \nabla \Phi \quad (4.42)$$

Similar operators can be constructed to generate rotations and Lorentz transformations, see the book by Itzykson & Zuber.

We finally note that the following particle number operator commutes with the Hamiltonian:

$$N = \sum_{\mathbf{k} \in \text{BZ}} a_{\mathbf{k}}^\dagger a_{\mathbf{k}} \quad (4.43)$$

It is curious to note that this operator is of course positive definite. The problem with the ‘negative energy solutions’ of the classical theory apparently does not appear in the quantum theory: both total energy and particle number are positive. However, it is not easy to give a local formulation to the particle number (density): to ensure that this is positive as well, one typically needs some spatial smearing.

Field propagator and commutator

To finish the discussion of the free Klein-Gordon theory, let us work out the commutator of the field operator between two space-time points x and y . If $\Phi(x)$ and $\Phi(y)$ are ‘independent’ observables, this should be zero. Any non-zero result is indicating that the field operator at event x can influence the operator at y : this is of course a check of causality in the relativistic setting. Using $[a_{\mathbf{k}}, a_{\mathbf{k}'}^\dagger] = \delta_{\mathbf{k}\mathbf{k}'}$, we get

$$i[\Phi(x), \Phi(y)] = \sum_{\mathbf{k} \in \text{BZ}} \frac{i}{2\omega_{\mathbf{k}}V} \left(e^{-ik_\mu(x^\mu - y^\mu)} - e^{ik_\mu(x^\mu - y^\mu)} \right) \quad (4.44)$$

where the relativistically invariant integration measure $d^3k/\omega_{\mathbf{k}}$ appears in the $V \rightarrow \infty$ limit (see Exercise 6.2)

$$i[\Phi(x), \Phi(y)] = \int \frac{d^3k}{(2\pi)^3\omega_{\mathbf{k}}} \sin k_\mu(x^\mu - y^\mu) \equiv D(x - y) \quad (4.45)$$

In Exercise 6.2, you have also shown that the function $D(x)$ only depends on the Minkowski distance $x_\mu x^\mu$. In particular, for $x^0 = 0$, one has $D(0, \mathbf{x}) = 0$ because the function under the integral is odd in \mathbf{k} . By Lorentz invariance, this property holds for any space-like separation: we find that indeed, the field operators commute if the two events are not inside each other’s light cone.

For time-like separations, one can analyze $D(t, \mathbf{0})$ as a function of t . With the change of variable $k \mapsto \omega_{\mathbf{k}}$, one can show that

$$D(t, \mathbf{0}) = -\frac{m}{8\pi|t|} J_1(mt) \quad (4.46)$$

where J_1 is the first-order Bessel function. The function (4.46) jumps at $t = 0$ from positive to negative values (the size of the jump is $-m^2/8\pi^3$) and decays smoothly to zero on a time scale $1/m$ with oscillations that have an envelope $\propto t^{-3/2}$. This implies that the commutator is nonzero on the light cone. Nonzero values are found also in the interior of the light cone (in the ‘forward’ and ‘backward’ directions) with a width of order $1/m$.

The commutator $D(x - y) = i[\Phi(x), \Phi(y)]$ is thus a c-number valued distribution; it is called the ‘propagator’ of the quantized Klein-Gordon field. This special propagator is not ‘causal’ in the sense that it propagates the field into the future (then it would be zero on one side of the light cone), but rather contains a superposition of incoming and outgoing spherical waves.

4.2.3 Application: particle decay

A key fact that must be implemented by quantum field theory is the creation and annihilation of particles. We have already seen that the operators a^\dagger and a with the same name can do this job. But as we want to implement the ‘conversion’ of particles of one species into another one, we have to specify an interaction between quantum fields. On the level of charged fields coupled to the electromagnetic field, the interaction Hamiltonian can be found from the minimal coupling procedure, but one has to quantize the electromagnetic field

⁵One can actually show that $\partial_t D(0, \mathbf{x}) = -\delta(\mathbf{x})$.

as well which comes with some technical difficulties. We discuss here a simpler example: a ‘self-interacting Klein-Gordon field’, also known as Φ^3 -theory.

In the phonon picture, this models the anharmonicity of the crystal lattice. As a consequence, the phonon modes are no longer normal modes of the system, but mode coupling and mode conversion occurs. In the quasi-particle language: a given phonon is unstable and decays into a pair of phonons.

To describe the decay of particles in quantum field theory, we need two ingredients: • Fermi’s Golden Rule, as coming from time-dependent perturbation theory, and • an interaction Hamiltonian that is not just bilinear in the field operators.

Fermi’s Golden Rule

This is a key result of time-dependent perturbation theory. See Appendix 5 for details. The result is the following transition probability per unit time for an initial state $|i\rangle$, summed over all final states $|f\rangle$ allowed by energy conservation:

$$\gamma_{i \rightarrow f} = \frac{dw_{i \rightarrow f}}{dT} = \frac{2\pi}{\hbar} \sum_f |(f|V|i)|^2 \delta(E_f - E_i) \quad (4.47)$$

Here, E_i and E_f are the initial and final energies, and V is the interaction Hamiltonian.

Interaction potential. We take here the simplest interaction within a scalar field theory:

$$V = \lambda \int_V d^3x \Phi^3(\mathbf{x}) \quad (4.48)$$

In the phonon picture, this corresponds to a nonlinear term in the vibrations of the lattice. In the field theory, it is called a ‘self-interaction’.

What kind of matrix elements involving this V can be nonzero? We shall focus on initial states containing one particle, $|i\rangle = a_{\mathbf{p}}^\dagger|\text{vac}\rangle$. By inserting the mode expansion (4.37) of the field operator, we get for example terms of the form

$$V \propto \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3} (\dots) a_{\mathbf{k}_1}^\dagger a_{\mathbf{k}_2}^\dagger a_{\mathbf{k}_3} \quad (4.49)$$

that create three particles. However, since this requires an energy of at least mc^2 for each particle, this process cannot conserve energy. One must change the energy of the initial particle, but this requires an operator sequence of the form $a_{\mathbf{k}}^\dagger a_{\mathbf{k}'}$ to be applied to $|i\rangle$.

Another group of terms is of the form

$$a_{\mathbf{k}_1} a_{\mathbf{k}_2}^\dagger a_{\mathbf{k}_3}$$

where the initial particle is annihilated (if $\mathbf{k}_3 = \mathbf{p}$). The vacuum expectation value of $a_{\mathbf{k}_1} a_{\mathbf{k}_2}^\dagger$ is nonzero if $\mathbf{k}_1 = \mathbf{k}_2$ so that it seems that the initial particle could simply disappear. But again, this process violates energy conservation because $E_i = E_{\mathbf{p}} = (m^2 + \mathbf{p}^2)^{1/2}$, while $E_f = 0$ (zero particles left).

We find a nonzero contribution for the decay into a pair of particles, coming from the combination

$$a_{\mathbf{k}_1}^\dagger a_{\mathbf{k}_2}^\dagger a_{\mathbf{k}_3} \quad (4.50)$$

provided $\mathbf{k}_3 = \mathbf{p}$. If we take as final state $\langle f | = \langle \text{vac} | a_{\mathbf{p}_1} a_{\mathbf{p}_2}$ we find a nonzero matrix element for the terms (4.50) with $\mathbf{k}_1 = \mathbf{p}_1$ and $\mathbf{k}_2 = \mathbf{p}_2$ (or vice versa). The space integral in Eq.(4.48) leads to the momentum conservation rule, $V \delta_{\mathbf{p}_1 + \mathbf{p}_2, \mathbf{p}}$, written here for discretized momenta in a finite normalization volume.

We finally integrate over the final momenta $\mathbf{p}_{1,2}$ of the two final particles to get the decay rate for the initial state. After some calculations, this leads to the integral

$$\gamma_{\mathbf{p}} = \pi \lambda^2 \int \frac{d^3 q}{(2\pi)^3} \frac{\delta(E_1(\mathbf{q}) + E_2(\mathbf{q}) - E_{\mathbf{p}})}{E_1(\mathbf{q}) E_2(\mathbf{q}) E_{\mathbf{p}}} \quad (4.51)$$

where $E_{\mathbf{p}}$ is again the energy of the initial particle. The momentum $\mathbf{q} = \mathbf{p}_1 - \mathbf{p}_2$ is the relative momentum of the particle pair. Its total momentum must be equal to \mathbf{p} . Therefore, the energies are

$$E_{1,2}(\mathbf{q})^2 = m^2 + \frac{1}{4}(\mathbf{p} \pm \mathbf{q})^2 \quad (4.52)$$

and the factors $1/E_{\mathbf{p}_i}$ come from the Lorentz-invariant integration measure in three-dimensional momentum space.

There is a threshold energy for the particle decay: since $E_1(\mathbf{q}) + E_2(\mathbf{q}) \geq 2m$, the initial particle must have an energy of at least $E_{\mathbf{p}} \geq 2m$ to decay into this pair. The maximum value of q that can contribute to the integral (4.51) is also easy to find out. An explicit evaluation of the integral is probably possible, but tedious. This kind of integrals is typical for results in elementary particle physics. One is saying that the decay rate is proportional to the ‘phase space’ available for the products of the decay (given conservation of total energy and momentum). On the other hand, the decay rate is proportional to the squared coupling constant λ^2 . This is typical for calculation based on Fermi’s Golden Rule (4.47).

4.3 Fermions

We would like to illustrate here a key point of quantum field theory: particles with spin $1/2$ (or spin $3/2, \dots$) have to be quantized in a different way from spin $0, 1, \dots$ particles. The latter ones (like phonons) are called ‘bosons’: they can occupy a given field mode in arbitrary numbers $a^\dagger a = 0, 1, 2, \dots$, as we have seen for phonons. This is forbidden for half-integer spins or ‘fermions’ where Pauli’s exclusion principle holds. There, $a^\dagger a = 0, 1$ are the only allowed occupation numbers for a given mode (this includes the spin quantum number, of course). We shall see that this is related to the occurrence of creation and annihilation operators that are ‘anti-commuting’. Our example is the Dirac field, but without the coupling to the electromagnetic field.

4.3.1 Anticommutators

We follow the quantization scheme that could be called ‘correspondence’ principle and take the classical energy and momentum density that follow with

Noether’s theorem from the Dirac Lagrangian. In the energy density integrated over all space, we promote the Dirac field to an operator in such a way that the energy operator generates time translations. Similarly for the momentum operator that generates space translations.

A reminder of Noether’s theorem: if a Lagrangian depends on the coordinate $x = (t, \mathbf{x})$ only via the fields Ψ (and their derivatives), the action integral is not changed under a translation of the field arguments (just shift the variables for the integration $d^4 x = dt d^3 x$). For an infinitesimal x -dependent translation by $\delta a^\mu(x)$, we have

$$\delta \Psi(x) = \delta a^\nu(x) \partial_\nu \Psi(x) \quad (4.53)$$

and the derivative of this quantity for $\partial_\mu \delta \Psi$. A calculation leads to

$$0 = \int d^4 x (\partial_\mu T^{\mu\nu}) \delta a_\nu(x) \quad (4.54)$$

$$T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \Psi_\alpha)} \partial^\nu \Psi_\alpha(x) - g^{\mu\nu} \mathcal{L} \quad (4.55)$$

The index α is summing over the components of the field, if needed. The quantity $T^{\mu\nu}$ is the energy-momentum tensor of the field theory. It is locally conserved, $\partial_\mu T^{\mu\nu} = 0$, which gives four conserved quantities.

For $\mu, \nu = 0, 0$, one gets the energy density whose integral is the total energy or Hamiltonian

$$H = \int d^3 x \left\{ \frac{\partial \mathcal{L}}{\partial(\partial_t \Psi_\alpha)} \partial_t \Psi_\alpha(x) - \mathcal{L} \right\} \quad (4.56)$$

we recognize the standard form here, since the first factor contains the momentum conjugate to $\Psi(x)$.

In the Dirac Lagrangian (2.38), the field $\Psi(x)$ and its conjugate $\Psi^\dagger(x)$ occur. We interpret these as different ‘components’ of the same field and get a total energy

$$H = \int d^3 x \frac{i}{2} (\Psi^\dagger \partial_t \Psi - \partial_t \Psi^\dagger \Psi) \quad (4.57)$$

The term $-\mathcal{L}$ does not contribute if we use the Dirac equation, restricting the analysis to field (operators) whose dynamics satisfies the Dirac equation.⁶ We expand the Dirac field into plane waves of particle-like and antiparticle-like character:

$$\Psi(x) = \sum_{\mathbf{k}, \sigma} \left[a_{\mathbf{k}\sigma} u_{\mathbf{k}\sigma} e^{-ik_\mu x^\mu} + b_{\mathbf{k}\sigma}^\dagger v_{\mathbf{k}\sigma} e^{ik_\mu x^\mu} \right] \quad (4.58)$$

where $u_{\mathbf{k}\sigma}$ and $v_{\mathbf{k}\sigma}$ are the spinor-valued amplitudes for a particle- and antiparticle-like plane-wave states with momentum \mathbf{k} and spin quantum number σ . More precisely: these spinors are those one gets if the Dirac solutions for particles at rest (positive or negative energies $\pm m$) are subject to a Lorentz transformation. This explains also the negative sign of the exponential $e^{ik_\mu x^\mu}$ in the antiparticle-like term. The operator $a_{\mathbf{k}\sigma}$ annihilates a particle, as we shall see below. The operator $b_{\mathbf{k}\sigma}^\dagger$ creates an anti-particle: we also see below that this choice is the one that ensures that the total energy is positive.

⁶In other words, $\mathcal{L} = 0$, for solutions of the equations of motion: this is the technical reason why the standard canonical quantization procedure runs into trouble.

By working out the total energy in terms of this plane wave expansion, we need some orthogonality relations that follow from the Lorentz invariance of $\bar{u}_{\mathbf{k}\sigma} v_{\mathbf{k}'\sigma'}$. More details are found in Itzykson & Zuber. The key observation is that the energy (4.57) brings, from the time derivative, the energies $\pm E_k = \pm(m^2 + k^2)^{1/2}$ into the game. The resulting total energy is

$$H = \sum_{\mathbf{k},\sigma} E_k \left[a_{\mathbf{k}\sigma}^\dagger a_{\mathbf{k}\sigma} - b_{\mathbf{k}\sigma} b_{\mathbf{k}\sigma}^\dagger \right] \quad (4.59)$$

that we want to interpret as a Hamiltonian. Let us proceed as usual first: the negative sign in front of the anti-particle like number operator $b_{\mathbf{k}\sigma}^\dagger b_{\mathbf{k}\sigma} = b_{\mathbf{k}\sigma} b_{\mathbf{k}\sigma}^\dagger - 1$ is coming from the fact that these modes have 'negative energy'. If we permit that this number operator has arbitrarily large eigenvalues, then the energy (4.59) is getting arbitrary large. This is what we want to avoid in order to get a (dynamically and thermodynamically) stable ground state.

The genial idea to make H positive is the rule

$$b_{\mathbf{k}\sigma} b_{\mathbf{k}\sigma}^\dagger = 1 - b_{\mathbf{k}\sigma}^\dagger b_{\mathbf{k}\sigma} \quad (4.60)$$

so that the number operator $b_{\mathbf{k}\sigma}^\dagger b_{\mathbf{k}\sigma}$ now appears with the opposite sign in Eq.(4.59). We can now identify the vacuum state as usual by $a_{\mathbf{k}\sigma}|vac\rangle = b_{\mathbf{k}\sigma}|vac\rangle = 0$. The (infinite) value of the energy in the vacuum state, $-\sum_{\mathbf{k},\sigma} E_k$, is subtracted (this looks like the energy when all modes with negative energy are filled with exactly one particle). After normal ordering, we get the following, nicely looking positive Hamiltonian

$$:H: = \sum_{\mathbf{k},\sigma} E_k \left[a_{\mathbf{k}\sigma}^\dagger a_{\mathbf{k}\sigma} + b_{\mathbf{k}\sigma}^\dagger b_{\mathbf{k}\sigma} \right]. \quad (4.61)$$

This prescription can be generalized by imposing that the particle and anti-particle creation and annihilation operators are anti-commuting:

$$\delta_{\mathbf{k}\sigma,\mathbf{k}'\sigma'} = \left\{ a_{\mathbf{k}\sigma}, a_{\mathbf{k}'\sigma'}^\dagger \right\} \equiv a_{\mathbf{k}\sigma} a_{\mathbf{k}'\sigma'}^\dagger + a_{\mathbf{k}'\sigma'}^\dagger a_{\mathbf{k}\sigma}, \quad (4.62)$$

similarly for the $b_{\mathbf{k}\sigma}$, and

$$0 = \left\{ a_{\mathbf{k}\sigma}, a_{\mathbf{k}'\sigma'} \right\} = \left\{ a_{\mathbf{k}\sigma}, b_{\mathbf{k}'\sigma'}^\dagger \right\} = \left\{ a_{\mathbf{k}\sigma}, b_{\mathbf{k}'\sigma'} \right\} \quad (4.63)$$

Note that the relations involving a and a are 'new': we did not have to specify this separately in the boson case. This implies in particular that when one applies a creation (or annihilation) operator twice, one gets nothing:

$$(a_{\mathbf{k}\sigma}^\dagger)^2 |vac\rangle = \frac{1}{2} \left\{ a_{\mathbf{k}\sigma}^\dagger, a_{\mathbf{k}\sigma}^\dagger \right\} |vac\rangle = 0 \quad (4.64)$$

It is not possible to create more than one particle in a given mode: this implements the Pauli exclusion principle.

Note, however, that the fermionic (anti-)commutation rules still permit the usual interpretation of the number operator. This comes from the following magic algebra:

$$\left[a_k^\dagger a_k, a_{k'}^\dagger \right] = a_k^\dagger \left\{ a_k, a_{k'}^\dagger \right\} - \left\{ a_k^\dagger, a_{k'}^\dagger \right\} a_k = a_k^\dagger \delta_{kk'} \quad (4.65)$$

so that the action of a_k^\dagger increases the particle number $a_k^\dagger a_k$ in mode k by one.

Similarly, after 'normal ordering', the total number of particles becomes

$$N = \int d^3x :\Psi^\dagger \Psi: = \sum_{\mathbf{k},\sigma} \left[a_{\mathbf{k}\sigma}^\dagger a_{\mathbf{k}\sigma} - b_{\mathbf{k}\sigma}^\dagger b_{\mathbf{k}\sigma} \right]. \quad (4.66)$$

This can now be negative or positive! It is funny that in the quantum field theory, this does no longer pose a problem.⁷ The negative particle number can be interpreted as the difference between 'particles' and 'anti-particles' or as the total charge (since particles and anti-particles have opposite charges).

To summarize, in the quantization of the Dirac field, one imposes that the total energy be positive (increasing from the vacuum state as particles are created), and this can be done by imposing anti-commutation rules for the mode operators $a_{\mathbf{k}\sigma}$ and $a_{\mathbf{k}\sigma}^\dagger$. These anti-commutation rules automatically imply the Pauli principle: a given mode for spin 1/2 particles can never be filled by more than one particle.

The filled Fermi sea. We could have formulated the Dirac quantization in terms of the operator $c_{\mathbf{k}\sigma} = b_{\mathbf{k}\sigma}^\dagger$. In that case, the energy for an anti-particle mode would have scaled like $-E_k c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma}$. We can re-phrase the preceding construction by saying that • the number operator involved here must have the maximum value 1; • the ground state is given by the 'Fermi sea', a state where all c -modes are filled,

$$|\text{Fermisea}\rangle = \prod_{\mathbf{k}\sigma} c_{\mathbf{k}\sigma}^\dagger |\text{empty}\rangle;$$

and • Pauli's principle holds, $(c_{\mathbf{k}\sigma}^\dagger)^2 = 0$. All these elements are automatically verified by using anti-commutators. In fact, when using anti-commutators, annihilation and creation operators can somehow be exchanged: 'create an anti-particle', b^\dagger , is like 'destroy a particle of the Fermi sea', c . On the two-dimensional Hilbert space spanned by the states $b^\dagger|vac\rangle = (1, 0)^T$ and $|vac\rangle = (0, 1)^T$, the operators b and b^\dagger are represented by the 'Pauli matrices'

$$b^\dagger \leftrightarrow \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad b \leftrightarrow \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

whose square is obviously zero.

4.3.2 Many-body theory

To conclude, we would like to illustrate the connection between the quantum field theory formulated up to now and the usual non-relativistic many-body quantum mechanics. You are probably familiar with the fact that for systems of indistinguishable particles, quantum statistics comes into play: the wave function $\psi(\mathbf{x}_1, \dots, \mathbf{x}_N)$ must be symmetric or anti-symmetric under particle exchange. For example,

$$\text{bosons/fermions: } \psi(\mathbf{x}_2, \mathbf{x}_1, \mathbf{x}_3, \dots, \mathbf{x}_N) = \pm \psi(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \dots, \mathbf{x}_N) \quad (4.67)$$

More generally the sign $\varepsilon(\pi)$ of the permutation π of particle coordinates occurs in $\psi(\mathbf{x}_{\pi 1}, \dots, \mathbf{x}_{\pi N})$ for fermions, while the many-boson wave function is totally symmetric under permutations.

⁷In fact, the quantum-mechanical probability that must be positive in quantum theory, is 'living' in another space: it is related to the complex probability amplitudes in quantum state superpositions like $\alpha|vac\rangle + \beta a_k|vac\rangle$.

N -particle sectors

What is the connection to the field operator? The general picture is the following: the field operator $\Psi(\mathbf{x})$ connects parts of the Hilbert space that differ by one in the total particle number (it is a combination of operators that create or annihilate one particle). However, these 'sectors' typically evolve independently since the particle number N commutes with the Hamiltonian. In a simple non-relativistic theory, one has

$$H = \sum_k E_k a_k^\dagger a_k, \quad N = \sum_k a_k^\dagger a_k. \quad (4.68)$$

N commutes with H because one requires an energy mc^2 to create an extra particle. The 'one-particle sector' \mathcal{H}_1 is spanned by the states $a_k^\dagger |\text{vac}\rangle$. One can thus ask: what is the 'wave function' of these states? If the k label plane wave states, then $a_k^\dagger |\text{vac}\rangle$ is an eigenstate of the momentum operator \hat{P} with eigenvalue k :

$$\hat{P} a_k^\dagger |\text{vac}\rangle = \sum_k k a_k^\dagger a_k a_k^\dagger |\text{vac}\rangle = k a_k^\dagger |\text{vac}\rangle \quad (4.69)$$

In the non-relativistic limit, it is useful to separate the field operator into positive and negative frequency components Ψ and Ψ^\dagger . The positive frequency part contains only annihilation operators:⁸

$$\Psi(\mathbf{x}, t) = \sum_k a_k(t) \phi_k(\mathbf{x}), \quad \Psi^\dagger(\mathbf{x}, t) = [\Psi(\mathbf{x}, t)]^\dagger \quad (4.70)$$

with orthonormal mode functions $\phi_k(\mathbf{x})$.

One-particle states.

The field operator $\Psi^\dagger(\mathbf{x}')$, when acting on the vacuum state, produces a quantum state in the one-particle sector \mathcal{H}_1 . Its probability amplitude in the momentum representation is given by (up to a normalization)

$$\tilde{\psi}(k; \mathbf{x}') = \langle \text{vac} | a_k \Psi^\dagger(\mathbf{x}') | \text{vac} \rangle \quad (4.71)$$

With the mode expansion (4.70), adding (subtracting) the vanishing average $\langle \text{vac} | \Psi^\dagger(\mathbf{x}') a_k | \text{vac} \rangle$ and using the (anti)-commutation rules for the a_k 's, we get

$$\tilde{\psi}(k; \mathbf{x}') = \phi_k^*(\mathbf{x}') \quad (4.72)$$

In the position representation, we thus have a 'wave function'

$$\psi(\mathbf{x}; \mathbf{x}') = \sum_k \phi_k^*(\mathbf{x}') \phi_k(\mathbf{x}) = \delta(\mathbf{x} - \mathbf{x}') \quad (4.73)$$

where we used the completeness relation of the mode functions $\phi_k(\mathbf{x})$. Hence, $\Psi^\dagger(\mathbf{x}') |\text{vac}\rangle$ is a state localized at the position \mathbf{x}' , or in other words:

⁸In the Klein-Gordon theory, the separation into positive and negative frequencies is natural. In the case of the relativistic Dirac field $\Psi_D(x)$, one can write $\Psi_D(x) \approx e^{-imc^2 t/\hbar} \Psi(x)$ and make $\Psi(x)$ slowly varying in time by averaging it over a time scale of order \hbar/mc^2 . This makes the anti-particle contribution disappear that evolves at negative frequencies.

the field operator $\Psi^\dagger(\mathbf{x}')$ [$\Psi(\mathbf{x}')$] creates [annihilates] a particle at the position \mathbf{x}' .

We can interpret the same calculation in terms of the wave function (in the usual sense, i.e., in the position representation) of the states created by a_k^\dagger . Projecting onto the position eigenstate $\Psi^\dagger(\mathbf{x}) |\text{vac}\rangle$, we have

$$\langle \text{vac} | \Psi(\mathbf{x}) a_k^\dagger | \text{vac} \rangle = \phi_k(\mathbf{x}) \quad (4.74)$$

Hence the plane wave mode function is also the quantum-mechanical wave function for the one-particle state associated with the mode k .

Two-particle states.

Two-particle states are generated by the basis states $a_k^\dagger a_{k'}^\dagger |\text{vac}\rangle$ with $k \neq k'$ in the Fermion case. To compute their wave function $\psi(\mathbf{x}_1, \mathbf{x}_2)$, we project onto the two-particle position state $\Psi^\dagger(\mathbf{x}_2) \Psi^\dagger(\mathbf{x}_1) |\text{vac}\rangle$:

$$\psi(\mathbf{x}_1, \mathbf{x}_2) = \langle \text{vac} | \Psi(\mathbf{x}_1) \Psi(\mathbf{x}_2) a_k^\dagger a_{k'}^\dagger | \text{vac} \rangle \quad (4.75)$$

By a calculation similar to the previous one, we shift the annihilation operators in $\Psi(\mathbf{x}_1)$ and $\Psi(\mathbf{x}_2)$ through a_k^\dagger and $a_{k'}^\dagger$ and use the (anti)commutation rules. The result is

$$\psi(\mathbf{x}_1, \mathbf{x}_2) = \phi_{k'}(\mathbf{x}_1) \phi_k(\mathbf{x}_2) \pm \phi_k(\mathbf{x}_1) \phi_{k'}(\mathbf{x}_2) \quad (4.76)$$

which is even (odd) under the exchange of the coordinates \mathbf{x}_1 and \mathbf{x}_2 , depending on Bose or Fermi statistics. We could have anticipated this result: since the operators a_k^\dagger and $a_{k'}^\dagger$ (anti)commute, the two-particle state $a_k^\dagger a_{k'}^\dagger |\text{vac}\rangle$ must be even (odd) in the quantum numbers k and k' .

Similarly, one can show that the state $a_{k_1}^\dagger \dots a_{k_N}^\dagger |\text{vac}\rangle$ is totally (anti)symmetric in the quantum numbers k_1, \dots, k_N , and its wave function $\psi(\mathbf{x}_1, \dots, \mathbf{x}_N)$ behaves similarly.

Interacting particles in second quantization

To conclude, we write down the link between the usual formulation of a quantum many-body theory for particles that interact via a potential $V(\mathbf{r})$ and the quantum field theory (or 'second quantized') language. The many-body Hamiltonian is, for N indistinguishable particles

$$H = \sum_{\alpha=1}^N \left(\frac{\mathbf{p}_\alpha^2}{2m} + W(\mathbf{x}_\alpha) \right) + \frac{1}{2} \sum_{\alpha \neq \beta} V(\mathbf{x}_\alpha - \mathbf{x}_\beta) \quad (4.77)$$

where $W(\mathbf{x})$ is an 'external potential' that is the same for all particles. Of course, the momentum operator is $\mathbf{p}_\alpha = -i\hbar \nabla_\alpha$, the derivative with respect to the α 'th coordinate. The particle density, for example, is given by the operator

$$n(\mathbf{x}) = \sum_\alpha \delta(\mathbf{x} - \mathbf{x}_\alpha) \quad (4.78)$$

In second quantization, the particle density becomes

$$n(\mathbf{x}) = \Psi^\dagger(\mathbf{x}) \Psi(\mathbf{x}) \quad (4.79)$$

while the Hamiltonian is (in normally ordered form)

$$H = \int d^3x \left(\frac{\hbar^2}{2m} \nabla \Psi^\dagger \cdot \nabla \Psi + \Psi^\dagger(\mathbf{x}) W(\mathbf{x}) \Psi(\mathbf{x}) \right) + \frac{1}{2} \int d^3x d^3x' \Psi^\dagger(\mathbf{x}) \Psi^\dagger(\mathbf{x}') V(\mathbf{x} - \mathbf{x}') \Psi(\mathbf{x}') \Psi(\mathbf{x}). \quad (4.80)$$

This formulation is the starting point for the discussion of several phenomena in physics: electron waves in condensed matter and superconductivity, liquid Helium and superfluidity, atomic matter waves at low temperatures and Bose-Einstein condensation. Generally speaking, the temperature of the system is a key parameter here, and one is looking for equilibrium correlation functions of the form

$$\langle \Psi^\dagger(\mathbf{x}, t) \Psi(\mathbf{x}', t') \rangle_T, \quad \langle n(\mathbf{x}, t) n(\mathbf{x}', t') \rangle_T \quad (4.81)$$

These functions can actually be measured by interference or scattering experiments.

In some of these applications, one works in a statistical ensemble where the particle number N is not fixed because the system is in contact with a 'particle reservoir'. This can be a good approximation even in the case where one knows that N is conserved, because in the thermodynamic limit $N \rightarrow \infty$, the fluctuations in the particle number are getting small. (The Bose-Einstein condensation of an ideal gas shows a peculiar scaling with N in this respect, see M. Wilkens and C. Weiss, *J. mod. Optics* **44** (1997) 1801.) The ground state of the system in this ensemble is found by minimizing the so-called 'grand-canonical Hamiltonian'

$$K = H - \mu N \quad (4.82)$$

where μ is called the 'chemical potential'. Its value is fixed by the requirement that the average particle number coincides with a given number \bar{N} :

$$\bar{N} = \frac{\partial}{\partial \mu} \log \text{tr} \exp(-\beta K) \quad (4.83)$$

where the trace gives the partition function (*Zustandssumme*) in this ensemble.

Chapter 5

Extra: Zeitabhängige Störungstheorie

5.1 Problemstellung

Gegeben sei ein Hamiltonoperator

$$\hat{H}(t) = \hat{H}_0 + \hat{V}(t) \quad (5.1)$$

wobei die Zeitentwicklung unter \hat{H}_0 bekannt sei. Zum Zeitpunkt t_0 sei das System im Eigenzustand $|a\rangle$ von \hat{H}_0 präpariert. Gesucht ist die Wahrscheinlichkeit, das System zum Zeitpunkt t im Eigenzustand $|b\rangle$ zu finden:

$$w_{ab}(t) = |\langle b|U(t, t_0)|a\rangle|^2. \quad (5.2)$$

Hier ist $U(t, t_0)$ der Zeitentwicklungsoperator zum Hamiltonoperator (5.1). Dieser unitäre Operator ist schwieriger auszurechnen als der für das ungestörte System, und wir werden ihn als Potenzreihe im Potential $\hat{V}(t)$ suchen.

5.2 Vorarbeiten

5.2.1 Wiederholung: Eigenschaften des Zeitentwicklungsoperators

Bewegungsgleichung (= Schrödingergleichung)

$$i\hbar \frac{\partial}{\partial t} U(t, t_0) = \hat{H}(t) U(t, t_0) \quad (5.3)$$

Anfangsbedingung

$$U(t_0, t_0) = 1 \quad (5.4)$$

Differentialgleichung und Anfangsbedingung lassen sich kombinieren in Integralgleichung

$$U(t, t_0) = 1 + \frac{1}{i\hbar} \int_{t_0}^t d\tau \hat{H}(\tau) U(\tau, t_0) \quad (5.5)$$

Beweis: differenzieren; $t = t_0$ einsetzen.

Schwierigkeit: Gl.(5.5) ist eine Integralgleichung, in der $U(t, t_0)$ auf beiden Seiten auftritt.

5.2.2 Wechselwirkungsbild

Idee: zwar können wir den Zeitentwicklungsoperator zu $\hat{H}(t)$ nicht bestimmen, wohl aber den zu \hat{H}_0 :

$$U_0(t, t_0) = \exp\left(-i\hat{H}_0(t - t_0)/\hbar\right) \quad (5.6)$$

Seine Wirkung in der Energiedarstellung $|a\rangle, |b\rangle, \dots |m\rangle, \dots$ ist diagonal:

$$U_0(t, t_0)|m\rangle = e^{-iE_m(t-t_0)/\hbar}|m\rangle \quad (5.7)$$

Wir definieren einen neuen Zustand $|\tilde{\psi}(t)\rangle$ durch folgende Vereinbarung

$$|\psi(t)\rangle = U_0(t, t_0)|\tilde{\psi}(t)\rangle \quad (5.8)$$

Wozu ist das gut? Aus der Schrödingergleichung kann man folgende Bewegungsgleichung für $|\tilde{\psi}(t)\rangle$ ausrechnen

$$i\hbar \frac{\partial}{\partial t} |\tilde{\psi}(t)\rangle = \tilde{V}(t)|\tilde{\psi}(t)\rangle \quad (5.9)$$

wobei

$$\tilde{V}(t) = U_0^\dagger(t, t_0)V(t)U_0(t, t_0) \quad (5.10)$$

Vorteil von Gl.(5.9): die Zeitentwicklung hängt nur noch vom Wechselwirkungsoperator ab.

Wichtiger **Begriff**: mit Hilfe der unitären Transformation (5.8) begibt man sich ins **Wechselwirkungsbild**.

5.2.3 von Neumann-Reihe

Zeitentwicklungsoperator zur Bewegungsgleichung (5.9):

$$|\tilde{\psi}(t)\rangle = U_I(t, t_0)|\tilde{\psi}(t_0)\rangle \quad (5.11)$$

Anfangsbedingung wie vorher. Damit Integralgleichung:

$$U_I(t, t_0) = \mathbf{1} + \frac{1}{i\hbar} \int_{t_0}^t d\tau \tilde{V}(\tau)U_I(\tau, t_0) \quad (5.12)$$

Wir sind immer noch nicht viel weiter, wenn wir nicht diese Gleichung lösen. Wir bekommen aber eine Potenzreihenentwicklung (die "von Neumann-Reihe"), wenn wir (5.12) iterieren, d.h. auf der rechten Seite die linke Seite

einsetzen. Man erhält damit:

$$U_I(t, t_0) = \sum_{n=0}^{\infty} U_I^{(n)}(t, t_0) \quad (5.13)$$

$$U_I^{(0)}(t, t_0) = \mathbf{1} \quad (5.14)$$

$$U_I^{(1)}(t, t_0) = \frac{1}{i\hbar} \int_{t_0}^t d\tau \tilde{V}(\tau) \quad (5.15)$$

$$U_I^{(2)}(t, t_0) = -\frac{1}{\hbar^2} \int_{t_0}^t d\tau_1 \int_{\tau_1 > \tau_2}^t d\tau_2 \tilde{V}(\tau_1)\tilde{V}(\tau_2) \quad (5.16)$$

...

Vornehm kann man diese Reihe auch als "zeitgeordnete Exponentialfunktion" schreiben:

$$U_I(t, t_0) = \mathcal{T} \exp\left(\frac{1}{i\hbar} \int_{t_0}^t d\tau \tilde{V}(\tau)\right) \quad (5.17)$$

wobei \mathcal{T} den (Super-)Operator der Zeitordnung bedeutet: in der Entwicklung des nachfolgende Ausdrucks sind Produkte von Operatoren $\tilde{V}(\tau_1)\tilde{V}(\tau_2)$ derart zu ordnen, dass die zeitlich späteren zuerst kommen ($\tau_1 > \tau_2$).

Ein wichtiges Ergebnis ist der Ausdruck (5.15) für den Zeitentwicklungsoperator in erster Ordnung: er ist dann ein einfaches Integral über die Wechselwirkung.

5.3 Fermi's Goldene Regel

Wir kehren jetzt zu der ursprünglichen Fragestellung (Gl. 5.2) zurück. Die gesuchte Übergangswahrscheinlichkeit enthält das Matrixelement

$$A_{ab} = \langle b|U(t, t_0)|a\rangle = \langle b|U_0(t, t_0)U_I(t, t_0)|a\rangle \quad (5.18)$$

In erster Ordnung wird daraus (die Wirkung von U_0 auf die Zustände $|a, b\rangle$ ist ein Phasenfaktor!)

$$\begin{aligned} A_{ab}^{(1)} &= \frac{e^{-iE_b(t-t_0)/\hbar}}{i\hbar} \int_{t_0}^t d\tau e^{iE_b(\tau-t_0)/\hbar} \langle b|V(\tau)|a\rangle e^{-iE_a(\tau-t_0)/\hbar} \\ &= \frac{e^{-iE_b(t-t_0)/\hbar}}{i\hbar} \int_{t_0}^t d\tau e^{i\omega_{ba}(\tau-t_0)} \langle b|V(\tau)|a\rangle \end{aligned} \quad (5.19)$$

$$\hbar\omega_{ba} = E_b - E_a \quad \text{BOHR'SCHE ÜBERGANGSFREQUENZ} \quad (5.20)$$

Hier sehen wir, dass die Übergangswahrscheinlichkeit von einer Art Fouriertransformation des Wechselwirkungsoperators abhängt, wobei genau die Bohr'sche Übergangsfrequenz ins Spiel kommt.

Wir betrachten jetzt eine zeitunabhängige Störung:

$$\hat{V}(t) = V \quad (5.21)$$

Dann kann man das Zeitintegral leicht auswerten und findet

$$A_{ab}^{(1)} = \frac{1}{i\hbar} e^{i\omega_{ba}T/2} \langle b|V|a\rangle \frac{\sin(\omega_{ba}T/2)}{\omega_{ba}/2} \quad (5.22)$$

$$T = t - t_0 \quad \text{DAUER DER WECHSELWIRKUNG} \quad (5.23)$$

Die Funktion $\sin(\Delta\omega T/2)/(\Delta\omega/2)$, die hier auftritt, besitzt für $\Delta\omega = 0$, also für $E_b = E_a$ einen scharfen Peak der Breite $2\pi/T$ (siehe Abbildung). Für große




Figure 5.1: Die Funktion $\sin(\Delta\omega T/2)/(\Delta\omega/2)$. Gestrichelt die Einhüllende der Oszillationen.

$|\Delta\omega|$ beschreibt sie Oszillationen, deren Amplitude proportional zu $1/|\Delta\omega|$ abfällt.

Wir erhalten folgenden Ausdruck für die Übergangswahrscheinlichkeit:

$$w_{ab}(t) = \frac{1}{\hbar^2} | \langle b|V|a \rangle |^2 \frac{\sin^2(\omega_{ba}T/2)}{\omega_{ba}^2/4} \quad (5.24)$$

wobei die Breite des Peaks bei $E_b \approx E_a$ um so kleiner wird, je größer die Dauer der Wechselwirkung ist. Dies sollte aus der Fourier-Transformation bekannt sein: Wirkt die Störung nur kurz, liegt eigentlich ein zeitabhängiges Problem vor, in dem die Energie nicht strikt erhalten ist (Unschärferelation Zeit-Energie).

5.3.1 Resonanzfall

Als erstes Beispiel betrachten wir den Fall, dass die Zustände $|a\rangle$ und $|b\rangle$ diskret sind und exakt $E_b = E_a$ gilt. Die Übergangswahrscheinlichkeit (5.24) wird dann

$$w_{ab}(t) = \frac{1}{\hbar^2} | \langle b|V|a \rangle |^2 T^2 \quad (5.25)$$

Dieser Ausdruck wächst quadratisch mit der Zeit T an. Dieses Verhalten ist charakteristisch für den Resonanzfall. Man bezeichnet die Frequenz $\Omega = | \langle b|V|a \rangle |/\hbar$ als RABI-FREQUENZ. Sie gibt die Zeitskala an, auf der die Wahrscheinlichkeit zunimmt, das System im Zustand $|b\rangle$ zu finden. Sie werden in der Quantenoptik ein Modell kennenlernen, in man sich auf zwei Zustände in einem Atom beschränkt (das "Zwei-Niveau-Atom"). Im Resonanzfall verhalten sich die Wahrscheinlichkeiten $w_a(t)$ und $w_b(t)$ wie $\cos^2(\Omega T)$ und $\sin^2(\Omega T)$. Für eine genügend kleine Rabi-Frequenz findet man den ersten Term der Näherung (5.25) wieder.

5.3.2 Anregung in ein Kontinuum

Als zweites Beispiel kommen wir zu Fermi's Goldener Regel, wie sie in den Büchern angegeben wird. Dazu betrachten wir den Fall, dass das Sys-

tem in ein Kontinuum $|B\rangle$ von Endzuständen in der Nähe des typischen Zustands $|b\rangle$ angeregt wird. Die relevante Größe ist nun die Summe der Übergangswahrscheinlichkeit $w_{ab}(t)$ über die möglichen Endzustände. Diese Summe kann als ein Integral über die Endenergie geschrieben werden

$$w_{a \rightarrow B}(t) = \int_{E_f - \varepsilon/2}^{E_f + \varepsilon/2} dE_b \rho(E_b) w_{ab}(t). \quad (5.26)$$

Dabei ist $\rho(E_b)$ die ZUSTANDSDICHTE im Energieraum: die Zahl $\rho(E)dE$ ist gleich der Anzahl der Zustände, deren Energie im Intervall $(E, E+dE)$ liegt. In der Wahrscheinlichkeit (5.26) werden alle Endzustände berücksichtigt, deren Energie bei $E_f = E_a$ liegt, mit einer Breite ε . Diese Breite gibt die "Energieunschärfe" des Detektors an.

Wir beschränken uns auf den Fall, dass die Energieunschärfe grösser als die Unschärfe aus der endlichen Wechselwirkungszeit ist:

$$\frac{2\pi\hbar}{T} \ll \varepsilon. \quad (5.27)$$

Aus dem Integral (5.26) erhalten wir:

$$w_{a \rightarrow B}(t) = \frac{1}{\hbar^2} \int_{E_f - \varepsilon/2}^{E_f + \varepsilon/2} dE_b \rho(E_b) | \langle b|V|a \rangle |^2 \frac{\sin^2(\omega_{ba}T/2)}{\omega_{ba}^2/4} \quad (5.28)$$

Die Funktion mit dem \sin enthält bekanntlich einen scharfen Peak an der Stelle $E_b = E_a$ mit der Breite $2\pi\hbar/T$. Um das Integral auszuwerten, nehmen wir weiter an, dass die Zustandsdichte und das Matrix-Element als Funktion von E_b auf der Skala ε langsam veränderlich sind. Damit können wir sie aus dem Integral herausziehen und erhalten:

$$\begin{aligned} w_{a \rightarrow B}(t) &= \frac{1}{\hbar^2} | \langle b|V|a \rangle |^2 \rho(E_f) \int_{E_f - \varepsilon/2}^{E_f + \varepsilon/2} dE_b \frac{\sin^2(\omega_{ba}T/2)}{\omega_{ba}^2/4} \\ &= \frac{2\pi T}{\hbar} | \langle b|V|a \rangle |^2 \rho(E_f) \end{aligned} \quad (5.29)$$

Im letzten Schritt haben wir ausgenutzt, dass für $E_f = E_a$ der gesamte Peak innerhalb der Energieunschärfe ε liegt und folgendes Integral benutzt:

$$\int_{-\infty}^{+\infty} dx \frac{\sin^2 x}{x^2} = \pi. \quad (5.30)$$

(Beweis: mit Hilfe des Residuensatzes.)

Unser Ergebnis (5.29) zeigt, dass die Übergangswahrscheinlichkeit in ein Kontinuum linear mit T ansteigt. Man beachte den Unterschied zum diskreten Endzustand, Gl.(5.25). Es bietet sich nun an, eine Übergangsrate einzuführen:

$$\gamma_{a \rightarrow B} = \frac{dw_{a \rightarrow B}}{dT} = \frac{2\pi}{\hbar} | \langle b|V|a \rangle |^2 \rho(E_f) \quad (5.31)$$

Dies ist nun endlich Fermi's Goldene Regel, wie sie in allen Büchern zur Quantenmechanik steht: die Übergangswahrscheinlichkeit in ein Kontinuum ist gleich dem Quadrat des Matrixelements der Störung, multipliziert mit der Zustandsdichte an der Energie des Endzustands.