

Chapter 1

The quantized field

We give an elementary introduction to the quantization of the electromagnetic field. We adopt the Coulomb gauge and a simple canonical framework. We then discuss examples for the quantum states of the field and the properties of vacuum fluctuations.

1.1 Canonical quantization

1.1.1 Fields

Maxwell equations. We want to quantize the vacuum Maxwell equations for the electromagnetic field. With a given charge and current density, these read

$$\begin{aligned} \nabla \cdot \mathbf{B} &= 0 & \varepsilon_0 \nabla \cdot \mathbf{E} &= \rho \\ \nabla \times \mathbf{E} + \partial_t \mathbf{B} &= 0 & \nabla \times \mathbf{B} - \frac{1}{c^2} \partial_t \mathbf{E} &= \mu_0 \mathbf{j} \end{aligned} \quad (1.1)$$

Introducing the scalar and vector potentials via

$$\mathbf{B} = \nabla \times \mathbf{A} \quad (1.2)$$

$$\mathbf{E} = -\nabla\phi - \partial_t \mathbf{A}, \quad (1.3)$$

the left column of (1.1) is identically fulfilled. The Coulomb law then becomes

$$-\varepsilon_0 \Delta\phi - \varepsilon_0 \nabla \partial_t \mathbf{A} = \rho \quad (1.4)$$

If we impose the Coulomb gauge, $\nabla \cdot \mathbf{A} = 0$, the vector potential drops out and the scalar potential is determined by the charge density alone:

$$-\varepsilon_0 \Delta \phi = \rho \quad (1.5)$$

In this gauge, the scalar potential is thus not a dynamical degree of freedom of the field: its dynamics is ‘enslaved’ by that of the charges. This holds with suitable boundary conditions such that the homogeneous equation $\Delta \phi = 0$ has no nontrivial solutions. In free space, with $\phi(\mathbf{x} \rightarrow \infty) \rightarrow 0$, we get

$$\phi(\mathbf{x}) = \frac{1}{4\pi\varepsilon_0} \int d^3x' \frac{\rho(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} \quad (1.6)$$

which is a superposition of well-known Coulomb potentials.

Wave equation. We are left with the wave equation for the vector potential

$$\nabla \times (\nabla \times \mathbf{A}) + \frac{1}{c^2} \partial_t^2 \mathbf{A} = \mu_0 (\mathbf{j} - \varepsilon_0 \partial_t \nabla \phi) \equiv \mu_0 \mathbf{j}_\perp. \quad (1.7)$$

On the right hand side, we have introduced the ‘transverse current’. Its divergence is zero because of Eq.(1.5) and charge conservation:

$$\nabla \cdot \mathbf{j}_\perp = \nabla \cdot \mathbf{j} + \partial_t \rho = 0. \quad (1.8)$$

The name ‘transverse’ comes from the fact that in spatial Fourier components, the current $\mathbf{j}_\perp(\mathbf{k})$ must be perpendicular to \mathbf{k} . One also says that the vector potential in the Coulomb gauge is transverse because $\nabla \cdot \mathbf{A} = 0$. The transversality of the source term in (1.7) ensures that if \mathbf{A} is transverse at one time, it is also transverse at all later times.

Conservation laws. Energy (Poynting theorem):

$$\partial_t u + \nabla \cdot \mathbf{S} = -\mathbf{j} \cdot \mathbf{E} \quad (1.9)$$

$$u = \frac{\varepsilon_0}{2} \mathbf{E}^2 + \frac{1}{2\mu_0} \mathbf{B}^2 \quad \text{energy density} \quad (1.10)$$

$$\mathbf{S} = \frac{1}{\mu_0} \mathbf{E} \times \mathbf{B} \quad \text{Poynting vector} \quad (1.11)$$

Momentum (see exercise). Charge: $\partial_t \rho + \nabla \cdot \mathbf{j} = 0$.

1.1.2 Matter

Let us consider that all matter is made from charged point particles with charges e_α and positions \mathbf{r}_α . The electric charge and current densities are then given by

$$\rho(\mathbf{x}, t) = \sum_{\alpha} e_{\alpha} \delta(\mathbf{x} - \mathbf{r}_{\alpha}(t)), \quad \mathbf{j}(\mathbf{x}, t) = \sum_{\alpha} e_{\alpha} \mathbf{v}_{\alpha}(t) \delta(\mathbf{x} - \mathbf{r}_{\alpha}(t)) \quad (1.12)$$

The sum runs over all the particles. Charge conservation is ensured provided that $\dot{\mathbf{r}}_{\alpha} = \mathbf{v}_{\alpha}$.

The point charges are thus the ‘sources’ for the electromagnetic field. But their motion is also influenced by the fields via the Newton-Lorentz equations:

$$\frac{d}{dt} \frac{m_{\alpha} \mathbf{v}_{\alpha}}{\sqrt{1 - v_{\alpha}^2/c^2}} = e_{\alpha} (\mathbf{E}(\mathbf{r}_{\alpha}) + \mathbf{v}_{\alpha} \times \mathbf{B}(\mathbf{r}_{\alpha})) \quad (1.13)$$

We use a relativistic framework here, and with the Lorentz factor, the time derivative is actually the one for the relativistic particle momentum. Note that in these equations, the coordinate \mathbf{r}_{α} enters generally in a nonlinear way. This differs from the Maxwell equations that are linear in the fields and potentials and whose solutions are linear in the charge and current distributions. (The superposition principle can be applied, see Eq.(1.6).) All nonlinear effects in optics can ultimately be traced back to the nonlinear response of matter to an applied electromagnetic field.

Quantization. The theory outlined so far describes physics at the end of the 19th century. It is unable to describe a stable state of matter because of the ‘radiation catastrophe’: positive and negative charges circle around each other, radiate electromagnetic waves and lose energy. Stable bound states exist only when the particle dynamics is quantized—recall the hydrogen atom.

Let us recall the typical energy and length scales for the hydrogen atom. The energy levels in Hydrogen are given by

$$E_n = -\frac{\text{Ryd}}{n^2} = -\frac{me^4}{2(4\pi\epsilon_0)^2\hbar^2} \frac{1}{n^2} \quad (1.14)$$

where the Rydberg constant is $1 \text{ Ryd} \approx 13.6 \text{ eV}$. (In cgs units, drop the factor $(4\pi\epsilon_0)^2$.) The size of the Hydrogen atom is of the order of the Bohr

radius

$$a_0 = \frac{4\pi\epsilon_0 \hbar^2}{me^2} \quad (1.15)$$

The typical wavelength of an electromagnetic wave resonant with a transition in Hydrogen is therefore of the order of

$$\lambda = \frac{\hbar c}{\text{Ryd}} = \frac{2}{\alpha_{\text{fs}}} a_0 \quad (1.16)$$

$$\frac{1}{\alpha_{\text{fs}}} = \frac{4\pi\epsilon_0 \hbar c}{e^2} \approx 137 \quad (1.17)$$

Here, α_{fs} is the fine structure constant. Its inverse can be understood as a measure of the speed of light in ‘atomic units’ (the natural units for the Hydrogen problem). The value $1/\alpha_{\text{fs}} \approx 137$ is fairly large. This means two things:

- the size of the hydrogen atom is small compared to the wavelength of resonant radiation: the Lorentz force (1.13) has therefore only a weak dependence on \mathbf{r}_α ;
- the typical velocity of an electron in the Hydrogen atom is in the non-relativistic regime: we can therefore use non-relativistic mechanics to describe the matter response.

This will justify several approximations for the atom-light interaction that we are going to make in the rest of the lecture.

1.1.3 Lagrange-Hamilton formulation

Lagrangian. We now proceed to quantize the wave equation (1.7) in the ‘canonical way’. The first step is to guess the corresponding Lagrangian. A good guess is the Lagrangian

$$L = - \sum_{\alpha} m_{\alpha} c^2 \sqrt{1 - \mathbf{r}_{\alpha}^2 / c^2} + \int d^3x \mathcal{L}_{F+I} \quad (1.18)$$

where the Lagrangian density for the field and its interaction with the electric charges is given by

$$\mathcal{L}_{F+I} = \frac{\epsilon_0}{2} (\dot{\mathbf{A}} + \nabla\phi)^2 - \frac{1}{2\mu_0} (\nabla \times \mathbf{A})^2 - \phi\rho + \mathbf{A} \cdot \mathbf{j}. \quad (1.19)$$

Via the Euler-Lagrange equations, one gets the Maxwell equations (1.5,1.7) for the fields and the Newton-Lorentz equation (1.13) for the particles. Note that the Lagrangian (1.18,1.19) is invariant under gauge transformations

$$\phi \mapsto \phi - \partial_t \chi, \quad \mathbf{A} \mapsto \mathbf{A} + \nabla \chi \quad (1.20)$$

where $\chi(\mathbf{x}, t)$ is an arbitrary smooth function. This gauge invariance is connected to charge conservation.

Coulomb gauge. We now proceed to specialize to the Coulomb gauge, simplify the Lagrangian and derive the Hamiltonian. We start with the terms involving the scalar potential in the Lagrangian (1.19). The mixed term is

$$\dot{\mathbf{A}} \cdot \nabla \phi = \nabla \cdot (\dot{\mathbf{A}} \phi) - \phi \nabla \cdot \dot{\mathbf{A}} \quad (1.21)$$

The first term is a divergence, and leads to a surface integral when integrated over the volume. We adopt the usual boundary condition that at the (infinitely remote) surface, the fields vanish: then this term is zero. The second term is zero in the Coulomb gauge $\nabla \cdot \mathbf{A} = 0$.

The term quadratic in the scalar potential is

$$(\nabla \phi)^2 = \nabla \cdot (\phi \nabla \phi) - \phi \Delta \phi = \nabla \cdot (\phi \nabla \phi) + \phi \rho / \epsilon_0 \quad (1.22)$$

using the Laplace equation (1.5). The second term thus combines with the interaction part $-\phi \rho$ in the Lagrangian that becomes $-\frac{1}{2} \phi \rho$. This energy can be interpreted as the Coulomb interaction energy between the charges:

$$V_{\text{Coul}} = \frac{1}{2} \int d^3x \phi \rho = \frac{1}{2} \int d^3x d^3x' \frac{\rho(\mathbf{x}) \rho(\mathbf{x}')}{4\pi \epsilon_0 |\mathbf{x} - \mathbf{x}'|} \quad (1.23)$$

$$= \frac{1}{2} \sum_{\alpha\beta} \frac{e_\alpha e_\beta}{4\pi \epsilon_0 |\mathbf{r}_\alpha - \mathbf{r}_\beta|} \quad (1.24)$$

where the factor $\frac{1}{2}$ ensures that all pairs of charges are only counted once. The divergent self-interaction for $\mathbf{r}_\alpha = \mathbf{r}_\beta$ also appears here. It is usually discarded. *The key point to note is that in the Coulomb gauge, the contribution of the scalar potential depends only on the particle coordinates. It is not a proper degree of freedom of the fields.*

To summarize, in the Coulomb gauge, the Lagrangian can be split into the following form:

$$L = - \sum_{\alpha} m_{\alpha} c^2 \sqrt{1 - v_{\alpha}^2/c^2} - V_{\text{Coul}}(\{\mathbf{r}_{\alpha}\}) + \int d^3x \mathcal{L}_{\text{F+I}}^{\perp} \quad (1.25)$$

with

$$\mathcal{L}_{\text{F+I}}^{\perp} = \frac{\varepsilon_0}{2} \dot{\mathbf{A}}^2 - \frac{1}{2\mu_0} (\nabla \times \mathbf{A})^2 + \mathbf{A} \cdot \mathbf{j}. \quad (1.26)$$

We have added the subscript \perp to remind ourselves that this is only valid if the vector potential is transverse.

Hamiltonian. For the Hamiltonian, we need the canonical momenta conjugate to \mathbf{r}_{α} and \mathbf{A} :

$$\mathbf{p}_{\alpha} = \frac{\partial L}{\partial \dot{\mathbf{r}}_{\alpha}} = \frac{m \dot{\mathbf{r}}_{\alpha}}{\sqrt{1 - \dot{\mathbf{r}}_{\alpha}^2/c^2}} + e_{\alpha} \mathbf{A}(\mathbf{r}_{\alpha}) \quad (1.27)$$

$$\mathbf{\Pi}(\mathbf{x}) = \frac{\delta L}{\delta \dot{\mathbf{A}}(\mathbf{x})} = \varepsilon_0 \dot{\mathbf{A}}(\mathbf{x}) \quad (1.28)$$

The particle momentum contains the relativistic kinetic momentum and an electromagnetic contribution. For the field momentum, it looks as if ε_0 where the ‘mass’ and $\dot{\mathbf{A}}$ the velocity.

Functional derivative. In the field case, we have a continuous collection of degrees of freedom, labelled by the space-points \mathbf{x} . This requires a generalization of the notion of a derivative to the infinite-dimensional case: the ‘functional derivative’ $\delta L/\delta \dot{\mathbf{A}}(\mathbf{x})$. It is the generalization of a gradient.

In mathematical terms, if we have a ‘functional’ $L[\mathbf{A}(\mathbf{x})]$, i.e. a mapping from the space of vector fields into the real numbers, its functional derivative is defined by the following approximation:

$$L[\mathbf{A}(\mathbf{x}) + \delta \mathbf{A}(\mathbf{x})] \approx L[\mathbf{A}(\mathbf{x})] + \int d^3x \left. \frac{\delta L}{\delta \mathbf{A}(\mathbf{x})} \right|_{\mathbf{A}(\mathbf{x})} \delta \mathbf{A}(\mathbf{x}) + \mathcal{O}(\delta \mathbf{A}^2) \quad (1.29)$$

Here, the second line is an example of a linear functional because the integral is linear in $\delta \mathbf{A}(\mathbf{x})$. The function with which the small deviation $\delta \mathbf{A}(\mathbf{x})$

is weighted under the integral *defines* the functional derivative. A mathematical theorem ensures that in a suitable space of functions, all linear functionals take this integral form.

It is a simple exercise to derive with (1.29) expression (1.28) for the field momentum $\mathbf{\Pi}$. We shall return to a less trivial example below.

The Hamiltonian is given by

$$H = \sum_{\alpha} \mathbf{p}_{\alpha} \cdot \dot{\mathbf{r}}_{\alpha} + \int d^3x \mathbf{\Pi} \cdot \dot{\mathbf{A}} - L \quad (1.30)$$

where the field part is the obvious generalization to a continuous set of degrees of freedom. Putting everything together, we get

$$H = \sqrt{[\mathbf{p}_{\alpha} - e_{\alpha} \mathbf{A}(\mathbf{r}_{\alpha})]^2 c^2 + m^2 c^4} + V_{\text{Coul}}(\{\mathbf{r}_{\alpha}\}) + \int d^3x \left[\frac{\mathbf{\Pi}^2}{2\epsilon_0} + \frac{(\nabla \times \mathbf{A})^2}{2\mu_0} \right] \quad (1.31)$$

where the interaction between matter and (transverse) field arises due to the ‘minimal coupling’ prescription only. (The term linear in \mathbf{j} in the Lagrangian cancels with the term linear in $\dot{\mathbf{r}}_{\alpha}$ in (1.30).)

Canonical equations. The motion of particles and fields in the Hamiltonian formalism can be described in a compact way in terms of Poisson brackets. We discuss this in some detail because they provide another example of functional derivatives and because they bear strong similarities to the commutators of the quantum theory. In addition, it turns out to be tricky to get the transverse wave equation (1.7).

The Poisson bracket provides the time evolution of any function Q (or functional) of the coordinates and momenta by

$$\dot{Q} = \{H, Q\} \quad (1.32)$$

where we define

$$\{A, B\} = \sum_{\alpha} \frac{\partial A}{\partial \mathbf{p}_{\alpha}} \cdot \frac{\partial B}{\partial \mathbf{r}_{\alpha}} - \frac{\partial A}{\partial \mathbf{r}_{\alpha}} \cdot \frac{\partial B}{\partial \mathbf{p}_{\alpha}} + \int d^3x \left[\frac{\delta A}{\delta \mathbf{\Pi}(\mathbf{x})} \cdot \frac{\delta B}{\delta \mathbf{A}(\mathbf{x})} - \frac{\delta A}{\delta \mathbf{A}(\mathbf{x})} \cdot \frac{\delta B}{\delta \mathbf{\Pi}(\mathbf{x})} \right] \quad (1.33)$$

Here, functional derivatives with respect to \mathbf{A} and $\mathbf{\Pi}$ appear for the fields. The Poisson bracket is antisymmetric in A and B and satisfies a ‘Jacobi identity’ (as does the commutator).

By working out $\{H, \mathbf{r}\}$, only the second term of the first line in (1.33) contributes, and we get (after some calculations) the relativistic relation between velocity and momentum, Eq.(1.27). This is left as an exercise. Similarly, one gets $\dot{\mathbf{A}} = \mathbf{\Pi}/\varepsilon_0$.

A more complicated calculation is needed for $\dot{\mathbf{\Pi}}$ where we have to evaluate

$$-\frac{\delta}{\delta \mathbf{A}(\mathbf{x})} \sqrt{[\mathbf{p}_\alpha - e_\alpha \mathbf{A}(\mathbf{r}_\alpha)]^2 c^2 + m^2 c^4} \quad (1.34)$$

We note first that

$$\frac{\delta \mathbf{A}(\mathbf{r}_\alpha)}{\delta \mathbf{A}(\mathbf{x})} = \delta(\mathbf{r}_\alpha - \mathbf{x}) \quad (1.35)$$

because the ‘evaluation functional’ $\mathbf{A} \mapsto \mathbf{A}(\mathbf{r}_\alpha)$ is of course a linear functional. (For mathematicians, this property *defines* the δ -function.) To proceed, we use the usual rules of differential calculus and get

$$-\frac{\delta}{\delta \mathbf{A}(\mathbf{x})} \sqrt{[\mathbf{p}_\alpha - e_\alpha \mathbf{A}(\mathbf{r}_\alpha)]^2 c^2 + m^2 c^4} = \delta(\mathbf{r}_\alpha - \mathbf{x}) \frac{e_\alpha (\mathbf{p}_\alpha - e_\alpha \mathbf{A}(\mathbf{r}_\alpha)) c^2}{\sqrt{[\mathbf{p}_\alpha - e_\alpha \mathbf{A}(\mathbf{r}_\alpha)]^2 c^2 + m^2 c^4}} \quad (1.36)$$

Performing the same calculations as for the particles’ equations of motion, this can be written as the current density $\mathbf{j}(\mathbf{x})$.

The last term now involves the derivative

$$-\frac{\delta}{\delta \mathbf{A}(\mathbf{x})} \frac{1}{2\mu_0} \int d^3x (\nabla \times \mathbf{A})^2 \quad (1.37)$$

that we handle with the mathematical definition (1.29). Consider a small change $\mathbf{a}(\mathbf{x})$ of the vector potential. To linear order, this gives a change

$$\begin{aligned} & (\nabla \times (\mathbf{A} + \mathbf{a}))^2 - (\nabla \times \mathbf{A})^2 \approx 2(\nabla \times \mathbf{a}) \cdot (\nabla \times \mathbf{A}) \\ & = 2\nabla \cdot [\mathbf{a} \times (\nabla \times \mathbf{A})] + 2\mathbf{a} \cdot [\nabla \times (\nabla \times \mathbf{A})] \end{aligned} \quad (1.38)$$

the first term is a divergence and vanishes after integrating over all space. The second one contains a multiplied with a weighting function so that we get

$$-\frac{1}{2\mu_0} \frac{\delta}{\delta \mathbf{A}(\mathbf{x})} \int d^3x (\nabla \times \mathbf{A})^2 = -\frac{1}{\mu_0} \nabla \times (\nabla \times \mathbf{A}) \quad (1.39)$$

Putting everything together, we have for the equation of motion of the vector potential:

$$\dot{\mathbf{\Pi}} = \varepsilon_0 \ddot{\mathbf{A}} = \mathbf{j}(\mathbf{x}) - \frac{1}{\mu_0} \nabla \times (\nabla \times \mathbf{A}) \quad \dots \text{wrong} \quad (1.40)$$

which is nearly equivalent to the wave equation (1.7). The point is that the source term is the ‘full current’, not its transverse part. This is actually an error in our calculation because we did not take into account the fact that the vector potential is restricted to be transverse.

A simple way to repair this is to use \mathbf{A}_\perp in the Lagrangian (1.25) and the Hamiltonian (1.31). Now, the link between \mathbf{A}_\perp and the ‘full’ \mathbf{A} is a linear functional (actually, a linear projector). This can be seen in the following way: consider an arbitrary \mathbf{A} and perform a gauge transformation (1.20) to remove the nonzero divergence. This fixes the ‘gauge function’ to satisfy

$$\Delta \chi = -\nabla \cdot \mathbf{A} \quad (1.41)$$

whose solution (vanishing at infinity) is given by a ‘superposition of Coulomb potentials’:

$$\chi(\mathbf{x}) = \frac{1}{4\pi} \int d^3x' \frac{\nabla' \cdot \mathbf{A}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} \quad (1.42)$$

where ∇' means the gradient with respect to \mathbf{x}' . After the gauge transformation, the now transverse vector potential is given by

$$\mathbf{A}_\perp(\mathbf{x}) = \mathbf{A}(\mathbf{x}) + \frac{1}{4\pi} \int d^3x' [\nabla' \cdot \mathbf{A}(\mathbf{x}')] \nabla \frac{1}{|\mathbf{x} - \mathbf{x}'|} \quad (1.43)$$

Now, $\nabla(1/r) = -\nabla'(1/r)$, and after one integration by parts we have

$$A_{\perp i}(\mathbf{x}) = \int d^3x' \delta_{ij}^\perp(\mathbf{x} - \mathbf{x}') A_j(\mathbf{x}') \quad (1.44)$$

where the so-called ‘transverse δ -function’ is given by

$$\delta_{ij}^\perp(\mathbf{x} - \mathbf{x}') = \delta_{ij} \delta(\mathbf{x} - \mathbf{x}') + \frac{1}{4\pi} \partial_j' \partial_i' \frac{1}{|\mathbf{x} - \mathbf{x}'|} \quad (1.45)$$

A careful evaluation of the second derivatives yields the explicit result

$$\delta_{ij}^\perp(\mathbf{x} - \mathbf{x}') = \frac{2}{3} \delta_{ij} \delta(\mathbf{x} - \mathbf{x}') - \frac{1}{4\pi} \left(\frac{\delta_{ij}}{r^3} - 3 \frac{r_i r_j}{r^5} \right), \quad \mathbf{r} = \mathbf{x} - \mathbf{x}' \quad (1.46)$$

By construction, the transverse δ -function acts like a usual δ -function on vector fields that are already transverse. We can interpret it as the ‘unit operator’ in the space of transverse vector fields.

Finally, if we write \mathbf{A}_\perp in the Hamiltonian, the equations of motion for $A_{\perp i}(\mathbf{x})$ lead to the following term

$$\int d^3x' \frac{\delta H}{\delta \Pi_j(\mathbf{x}')} \frac{\delta A_{\perp i}(\mathbf{x})}{\delta A_j(\mathbf{x}')} = \int d^3x' \frac{\delta H}{\delta \Pi_j(\mathbf{x}')} \delta_{ij}^\perp(\mathbf{x} - \mathbf{x}') = \left(\frac{\delta H}{\delta \Pi(\mathbf{x}')} \right)_{\perp i} \quad (1.47)$$

The functional derivative with respect to \mathbf{A} makes the transverse δ -function appear. The integral over \mathbf{x}' then projects the functional derivative with respect to Π into the transverse subspace. In this way, the time derivative of \mathbf{A}_\perp is transverse, as it should do in order to maintain the Coulomb gauge at all times.

The same procedure applies to the equation of motion for the field momentum: the bottomline is that the source current \mathbf{j} is ‘transversalized’, leading to the correct wave equation (1.7).

1.2 Canonical quantization

1.2.1 Mode expansion

The next step is to look for ‘normal modes’ of this field theory. We first identify some general requirements for the modes. The passage to plane waves is a little bit tricky and is treated in detail in the exercises. Note that we ignore for the moment the matter-field coupling: we focus on the field Hamiltonian only.

We adopt the expansion

$$\begin{pmatrix} \mathbf{A}(\mathbf{x}, t) \\ \Pi(\mathbf{x}, t) \end{pmatrix} = \sum_{\kappa} \mathbf{f}_{\kappa}(\mathbf{x}) \begin{pmatrix} q_{\kappa}(t) \\ p_{\kappa}(t) \end{pmatrix} \quad (1.48)$$

where the ‘mode functions’ $\mathbf{f}_{\kappa}(\mathbf{x})$ carry the space dependence and the ‘coordinates’ $q_{\kappa}(t)$ and ‘momenta’ $p_{\kappa}(t)$ the time-dependence.

The Coulomb gauge requires, of course, $\nabla \cdot \mathbf{f}_{\kappa}(\mathbf{x}) = 0$: the mode functions must be transverse.

We want the Hamiltonian to adopt a simple form using these modes. The space integral over Π^2 becomes simple if we impose the modes to be orthogonal:

$$\int d^3x \mathbf{f}_\kappa(\mathbf{x}) \cdot \mathbf{f}_{\kappa'}(\mathbf{x}) = N_\kappa \delta_{\kappa\kappa'} \quad (1.49)$$

where N_κ is a normalization constant that we fix later. The momentum-part of the Hamiltonian then becomes

$$\frac{1}{2\varepsilon_0} \int d^3x \mathbf{\Pi}^2 = \sum_\kappa N_\kappa \frac{p_\kappa^2}{2\varepsilon_0} \quad (1.50)$$

Similarly, for the integral over $(\nabla \times \mathbf{A})^2$. Integrating by parts:

$$\begin{aligned} & \int d^3x (\nabla \times \mathbf{f}_\kappa) \cdot (\nabla \times \mathbf{f}_{\kappa'}) \\ &= \int d\mathbf{A} \cdot [\mathbf{f}_\kappa \times (\nabla \times \mathbf{f}_{\kappa'})] + \int d^3x \mathbf{f}_\kappa \cdot [\nabla \times (\nabla \times \mathbf{f}_{\kappa'})] \end{aligned} \quad (1.51)$$

The volume integral, strictly speaking, must be evaluated over a finite volume only, otherwise, we could not work with a discrete set of mode labels κ . The boundary term can nevertheless be made to vanish if either (i) the mode function \mathbf{f}_κ or its curl $(\nabla \times \mathbf{f}_{\kappa'})$ is required to vanish on the boundary of the volume or (ii) periodic boundary conditions on ‘opposite faces’ of a cubic volume are assumed. The case (i) is appropriate for modes in a cavity with perfectly conducting boundaries: then, \mathbf{f}_κ is proportional to the electric field, and the integrand in Eq.(1.51) vanishes because the field is normal to the boundary. The case (ii) is the favorite one for theorists because the mode functions can be taken as plane waves. Note that the eigenfrequencies of the two cavities are not the same. We ignore for the moment the complications of complex mode functions (see details below and the exercises) and continue.

The volume integral in (1.51) is reduced to the orthogonality relation if we require the mode functions to be eigenfunctions of the (vector) Helmholtz equation:

$$\nabla \times (\nabla \times \mathbf{f}_\kappa) = \varepsilon_0 \mu_0 \omega_\kappa^2 \mathbf{f}_\kappa \quad (1.52)$$

(This equation is actually equivalent to the scalar Helmholtz equation for all components of \mathbf{f}_κ because of transversality.)

With all these assumptions taken together, the Hamiltonian for the field takes the form

$$H_F = \sum_{\kappa} \left[\frac{N_{\kappa}}{2\varepsilon_0} p_{\kappa}^2 + \frac{\varepsilon_0}{2} N_{\kappa} \omega_{\kappa}^2 q_{\kappa}^2 \right] \quad (1.53)$$

We now fix the normalization to be $N_{\kappa} = 1$ and get a sum of harmonic oscillator Hamiltonians, one for each mode κ with ‘mass’ ε_0 and frequency ω_{κ} .

Note: the construction of field modes is a ‘classical problem’ of electrodynamics, it has nothing to do with quantum mechanics. The word ‘quantization volume’ that is sometimes used (to ensure that the mode index κ is discrete) is therefore misleading. Quantization is something different, as we shall see now.

Mode operators. Quantization proceeds by promoting the p_{κ} and q_{κ} to operators with the commutation relations

$$\frac{i}{\hbar} [p_{\kappa}, q_{\kappa'}] = \delta_{\kappa\kappa'} \quad (1.54)$$

The choice for this commutator is similar to the one for the particle coordinates and momenta in ordinary quantum mechanics. So in the end, field quantization is nothing else but ordinary quantization, once the dynamics of the field is reduced to a discrete set of ‘normal modes’. The procedure that we have followed was first laid out by Dirac. It is called ‘canonical quantization’.

The commutator between the fields becomes

$$\frac{i}{\hbar} [\Pi_i(\mathbf{x}), A_j(\mathbf{x}')] = \sum_{\kappa} f_{\kappa i}(\mathbf{x}) f_{\kappa j}(\mathbf{x}') \stackrel{!}{=} \delta_{ij}^{\perp}(\mathbf{x} - \mathbf{x}') \quad (1.55)$$

The last equality is obtained by applying the canonical quantization scheme directly to the fields, care being taken that the fields (operators) ‘live’ in the space of transverse vector fields (operators). It means that the mode functions $f(\mathbf{x})$ form a *complete set* of functions in the transverse field space. To implement this equality, one takes in practice the limit of an infinitely large quantization volume where the sum degenerates into an integral. With discrete functions, one can actually represent only a ‘finite volume version’ of the transverse δ -function—the one that comes by applying the finite volume boundary conditions to the equation (1.41).

For the harmonic oscillator, creation and annihilation operators are a convenient tool to construct the Hilbert space of quantum states. In our context, these operators, a_κ^\dagger and a_κ , correspond to the ‘creation’ and ‘destruction’ of one ‘photon’. The mode coordinate and momentum operators are given by

$$q_\kappa = \sqrt{\frac{\hbar}{2\varepsilon_0\omega_\kappa}} (a_\kappa + a_\kappa^\dagger) \quad (1.56)$$

$$p_\kappa = \sqrt{\frac{\hbar\varepsilon_0\omega_\kappa}{2}} i (a_\kappa - a_\kappa^\dagger) \quad (1.57)$$

where we continue to write ε_0 for the oscillator mass and where the commutation relation is

$$[a_\kappa, a_{\kappa'}^\dagger] = \delta_{\kappa\kappa'} \quad (1.58)$$

The field Hamiltonian then takes the form

$$H_F = \sum_\kappa \frac{\hbar\omega_\kappa}{2} (a_\kappa a_\kappa^\dagger + a_\kappa^\dagger a_\kappa) = \sum_\kappa \hbar\omega_\kappa \left(a_\kappa^\dagger a_\kappa + \frac{1}{2} \right) \quad (1.59)$$

The last way of writing makes two essential things explicit:

- the energies (the energy eigenvalues!) of a given field mode are quantized in units of $\hbar\omega_\kappa$ (a ‘photon energy’) and the ‘number of photons’ is represented by the operator $a_\kappa^\dagger a_\kappa$.
- The ground state of the field corresponds to the state $|\text{vac}\rangle$ such that $a_\kappa|\text{vac}\rangle = 0$ for all κ . This is an energy eigenstate whose energy is infinity, $\frac{1}{2} \sum_\kappa \hbar\omega_\kappa$, the sum over the ‘zero-point energies’ of all the modes.

To summarize, we give the mode expansions of the vector potential that we have found

$$\mathbf{A}(\mathbf{x}, t) = \sum_\kappa \sqrt{\frac{\hbar}{2\varepsilon_0\omega_\kappa}} \mathbf{f}_\kappa(\mathbf{x}) \left(a_\kappa e^{-i\omega_\kappa t} + a_\kappa^\dagger e^{i\omega_\kappa t} \right). \quad (1.60)$$

We have used here the Heisenberg picture for the vector potential operator. From the Hamiltonian (1.59), it is easy to show that the operator $a_\kappa(t)$ evolves with a complex exponential $e^{-i\omega_\kappa t}$. This is also called a ‘positive frequency operator’. It can be shown that operators that destroy particles and lower the energy of a quantum state are always positive frequency operators.

Plane wave expansion. For completeness, we give here the plane-wave expansion for the field mode functions. These are complex, and therefore they are normalized according to

$$\int d^3x \mathbf{f}_\kappa^*(\mathbf{x}) \cdot \mathbf{f}_{\kappa'}(\mathbf{x}) = \delta_{\kappa\kappa'} \quad (1.61)$$

instead of Eq.(1.49). This can be ensured with the choice

$$\mathbf{f}_\kappa(\mathbf{x}) = \frac{1}{\sqrt{V}} \mathbf{u}_\kappa e^{i\mathbf{k}\cdot\mathbf{x}} \quad (1.62)$$

where V is the volume of the box with periodic boundary conditions, \mathbf{k} is a discrete wave vector: it increases in steps of $2\pi/V^{1/3}$ for a cubic box. And \mathbf{u}_κ is a ‘transverse’ polarization vector with the property $\mathbf{k} \cdot \mathbf{u}_\kappa = 0$. There are two mutually orthogonal choices of polarization for a given \mathbf{k} . (These can be complex, describing circular polarization.) The frequency of this mode is $\omega_\kappa = |\mathbf{k}|/\sqrt{\varepsilon_0\mu_0} = c|\mathbf{k}|$, as can be seen from the Helmholtz equation (1.52). The magnetic field is oriented along $\mathbf{k} \times \mathbf{u}_\kappa \equiv (\omega_\kappa/c)\mathbf{v}_\kappa$.

Finally, the quantized vector potential, electric and magnetic fields are given in terms of the following plane wave expansion

$$\mathbf{A}_\perp(\mathbf{x}, t) = \sum_\kappa \sqrt{\frac{\hbar}{2\varepsilon_0\omega_\kappa V}} \left(\mathbf{u}_\kappa e^{i(\mathbf{k}\cdot\mathbf{x} - \omega_\kappa t)} a_\kappa + \text{h.c.} \right). \quad (1.63)$$

$$\mathbf{E}_\perp(\mathbf{x}, t) = \sum_\kappa \sqrt{\frac{\hbar\omega_\kappa}{2\varepsilon_0 V}} \left(i\mathbf{u}_\kappa e^{i(\mathbf{k}\cdot\mathbf{x} - \omega_\kappa t)} a_\kappa + \text{h.c.} \right). \quad (1.64)$$

$$\mathbf{B}(\mathbf{x}, t) = \sum_\kappa \sqrt{\frac{\hbar\omega_\kappa\mu_0}{2V}} \left(i\mathbf{v}_\kappa e^{i(\mathbf{k}\cdot\mathbf{x} - \omega_\kappa t)} a_\kappa + \text{h.c.} \right). \quad (1.65)$$

Sometimes, you may encounter these formulas without the factor i in front (and with ‘+h.c.’). Then the operator $ia_{\mathbf{k}\lambda}$ is being used instead. Note that Eq.(1.64) gives only the ‘transverse’ part of the electric field. The ‘longitudinal’ part, $-\nabla\phi$, is determined according to (1.5) by the charge density.

Note: A useful shortcut to derive the prefactors is the following: for each mode, match the energy density $\frac{1}{2}\varepsilon_0\mathbf{E}^2 + (1/2\mu_0)\mathbf{B}^2$ to the photon energy per quantization volume, $(\hbar\omega_\kappa/V)(a_\kappa^\dagger a_\kappa + \frac{1}{2})$. In free space, the electric and magnetic energy densities are equal. Average over the spatial oscillations in \mathbf{E}^2 for simplicity.

Exercise. Show that the momentum of the field can be written as a sum over modes as well. With plane wave modes, the momentum per mode is quantized in units of $\hbar\mathbf{k}$, as expected. Arbitrary cavity modes involving sin or cos functions are not eigenfunctions of the momentum operators, therefore their momentum is not well-defined. A ‘cavity photon’ therefore does not have a well-defined momentum.

Exercise. Write the equation of motion for a mode operator $a_{\mathbf{k}}$ and include the source current. Solve it for known time-dependence of current.

1.2.2 Plane wave expansion

In many cases, plane waves are a more natural choice for the mode functions. But they are complex, and this leads to some technical difficulties. We outline below a quantization procedure that directly starts from plane waves.

Plane waves are an obvious choice in a box of volume V with periodic boundary conditions. We can expand

$$\mathbf{A}(\mathbf{x}, t) = \sum_{\mathbf{k}} \mathbf{A}_{\mathbf{k}}(t) e^{i\mathbf{k}\cdot\mathbf{x}} \quad (1.66)$$

Some simple manipulations then lead to the Lagrangian

$$L = \frac{\varepsilon_0 V}{2} \sum_{\mathbf{k}} \left(\dot{\mathbf{A}}_{\mathbf{k}} \cdot \dot{\mathbf{A}}_{-\mathbf{k}} - c^2 k^2 \mathbf{A}_{\mathbf{k}} \cdot \mathbf{A}_{-\mathbf{k}} + \mathbf{A}_{\mathbf{k}} \cdot \mathbf{j}_{-\mathbf{k}} \right). \quad (1.67)$$

This already decomposes into a sum over the wave vectors \mathbf{k} . Only waves with \mathbf{k} and $-\mathbf{k}$ are coupled. We can also decompose the complex vector in a basis of orthogonal polarization vectors (there are two of them with $\mathbf{k} \perp \mathbf{u}_{\mathbf{k}\lambda}$)

$$\mathbf{A}_{\mathbf{k}} = \sum_{\lambda} \mathbf{u}_{\mathbf{k}\lambda} A_{\mathbf{k}\lambda}. \quad (1.68)$$

For simplicity, we assume that the polarization vectors are real and $\mathbf{u}_{\mathbf{k}\lambda} = \mathbf{u}_{-\mathbf{k}\lambda}$. The complex number $A_{\mathbf{k}\lambda}$ still encodes two degrees of freedom that are related to the plane waves with wave vectors \mathbf{k} and $-\mathbf{k}$. This is apparent from the relation

$$\mathbf{A}_{\mathbf{k}}^* = \mathbf{A}_{-\mathbf{k}} \quad (1.69)$$

that follows from the fact that $\mathbf{A}(\mathbf{x}, t)$ is real. We now make the decomposition

$$A_{\mathbf{k}\lambda} = q_1 + iq_2, \quad A_{-\mathbf{k}\lambda} = q_1 - iq_2, \quad (1.70)$$

where the indices $\mathbf{k}\lambda$ have been dropped just for the clarity of presentation. In terms of these variables, the terms with \mathbf{k} and $-\mathbf{k}$ combine to give a sum of two harmonic oscillators:

$$L_{\mathbf{k}} = \varepsilon_0 V \left(\dot{q}_1^2 + \dot{q}_2^2 - c^2 k^2 (q_1^2 + q_2^2) + \text{interactions} \right) \quad (1.71)$$

(The factor $\frac{1}{2}$ drops out because both \mathbf{k} and $-\mathbf{k}$ give the same contribution.)

Quantization. Canonical quantization means that the coordinate q_1 and its canonically conjugate momentum

$$p_1 = \frac{\partial L}{\partial \dot{q}_1} = 2\varepsilon_0 V \dot{q}_1 \quad (1.72)$$

are replaced by operators with the commutator $[p_\alpha, q_\beta] = -i\hbar \delta_{\alpha\beta}$ ($\alpha, \beta = 1, 2$). The Hamiltonian is given by

$$H = \frac{1}{4\varepsilon_0 V} (p_1^2 + p_2^2) + \varepsilon_0 V c^2 k^2 (q_1^2 + q_2^2) + \text{interaction} \quad (1.73)$$

so that we are dealing with two uncoupled harmonic oscillators. We can use the well-known results from the Quantum Mechanics lecture if we write $\omega = ck$ and $m = 2\varepsilon_0 V$. Here ω is really a frequency (the light frequency for a plane wave), and m is not a mass because q is not a position coordinate, but has the dimensions of a vector potential.

The quantization of the harmonic oscillator is most easily written down in terms of the annihilation and creation operators a_α and a_α^\dagger . These are dimensionless quantities with commutator $[a_\alpha, a_\beta^\dagger] = \delta_{\alpha\beta}$. They allow to represent coordinates and momenta as

$$q_\alpha = \sqrt{\frac{\hbar}{4\varepsilon_0 V \omega}} (a_\alpha + a_\alpha^\dagger), \quad (1.74)$$

$$p_\alpha = -i\sqrt{\hbar\varepsilon_0 V \omega} (a_\alpha - a_\alpha^\dagger). \quad (1.75)$$

We thus get the following expansion of the vector potential

$$\begin{aligned} \mathbf{A}(\mathbf{x}) = \sum'_{\mathbf{k}, \lambda} \sqrt{\frac{\hbar}{4\varepsilon_0 V \omega_{\mathbf{k}}}} \mathbf{u}_{\mathbf{k}\lambda} \left\{ (a_1 + a_1^\dagger + i(a_2 + a_2^\dagger)) e^{i\mathbf{k}\cdot\mathbf{x}} \right. \\ \left. + (a_1 + a_1^\dagger - i(a_2 + a_2^\dagger)) e^{-i\mathbf{k}\cdot\mathbf{x}} \right\} \end{aligned} \quad (1.76)$$

The primed sum is to remind us that we sum only over ‘one half’ of k-space, regrouping \mathbf{k} and $-\mathbf{k}$. We now introduce the operators

$$b_1 = \frac{a_1 + ia_2}{\sqrt{2}}, \quad b_2 = \frac{a_1 - ia_2}{\sqrt{2}}. \quad (1.77)$$

They also satisfy the commutation relations $[b_\alpha, b_\beta^\dagger] = \delta_{\alpha\beta}$ and are therefore equally well suited to represent the normal modes of the field. They have the advantage that the plane wave expansion (1.76) can be written in the form

$$\mathbf{A}(\mathbf{x}) = \sum'_{\mathbf{k},\lambda} \sqrt{\frac{\hbar}{2\varepsilon_0 V \omega_{\mathbf{k}}}} \mathbf{u}_{\mathbf{k}\lambda} \{b_1 e^{i\mathbf{k}\cdot\mathbf{x}} + b_2 e^{-i\mathbf{k}\cdot\mathbf{x}} + \text{h.c.}\}. \quad (1.78)$$

We can now label the mode operators as $b_1 = a_{\mathbf{k}\lambda}$ and $b_2 = a_{-\mathbf{k}\lambda}$, so that the summation can be extended over all k-space:

$$\boxed{\mathbf{A}(\mathbf{x}) = \sum_{\mathbf{k},\lambda} \sqrt{\frac{\hbar}{2\varepsilon_0 V \omega_{\mathbf{k}}}} \{ \mathbf{u}_{\mathbf{k}\lambda} a_{\mathbf{k}\lambda} e^{i\mathbf{k}\cdot\mathbf{x}} + \text{h.c.} \}}. \quad (1.79)$$

This is the standard expansion of the vector potential in quantum electrodynamics.

Interaction and free field Hamiltonian. The interaction Hamiltonian with the transverse current takes the following form in the quantized theory:

$$H_{FA} = \int d^3x \mathbf{A}(\mathbf{x}) \cdot \mathbf{j}_\perp(\mathbf{x}, t) \quad (1.80)$$

$$= \sum_{\mathbf{k}\lambda} \sqrt{\frac{\hbar}{2\varepsilon_0 V \omega_{\mathbf{k}}}} \{ a_{\mathbf{k}\lambda} \mathbf{u}_{\mathbf{k}\lambda} \cdot \mathbf{j}_{\perp,\mathbf{k}}^*(t) + \text{h.c.} \}, \quad (1.81)$$

where $\mathbf{j}_{\perp,\mathbf{k}}(t)$ is the spatial Fourier transform.

The dynamics of the field itself is generated by the well-known collection of harmonic oscillators:

$$H_F = \sum_{\mathbf{k},\lambda} \hbar \omega_{\mathbf{k}} \left(a_{\mathbf{k}\lambda}^\dagger a_{\mathbf{k}\lambda} + \frac{1}{2} \right). \quad (1.82)$$

In the present context, we treated the current density as an ‘external’, given function with an explicit time-dependence. For this reason, there is no Hamiltonian required for it. If the motion of the charges is quantized as well, one needs, of course, an ‘atomic’ Hamiltonian to generate their dynamics.

Field commutators. From the mode expansions (1.79, 1.64, 1.65) and the commutation relations between the annihilation operators, one can compute the commutator between the electric and magnetic fields. In the Lagrangian description, one finds already that the vector potential and the electric field are (up to a factor $-\varepsilon_0$) canonically conjugate variables. This is only true in the space of ‘transverse vector functions’ (fields with zero divergence), however. In the quantized theory, the corresponding commutator is

$$[A_l(\mathbf{x}, t), E_m(\mathbf{x}', t)] = \frac{i\hbar}{\varepsilon_0} \delta_{lm}^\perp(\mathbf{x} - \mathbf{x}') \quad (1.83)$$

where δ_{lm}^\perp is the ‘transverse δ -function’ defined in Eq.(1.45). As discussed above, this distribution acts like a δ -function on fields with zero divergence, and projects an arbitrary vector field \mathbf{F} on its transverse part \mathbf{F}^\perp

$$F_l^\perp(\mathbf{x}) = \int d^3x' \delta_{lm}^\perp(\mathbf{x} - \mathbf{x}') F_m(\mathbf{x}') \quad (1.84)$$

By definition, the transverse part has zero divergence, $\nabla \cdot \mathbf{F}^\perp = 0$. The projection is most easily constructed in Fourier space

$$F_j^\perp(\mathbf{x}) = \int \frac{d^3k}{(2\pi)^3} \frac{k^2 \delta_{jl} - k_j k_l}{k^2} \tilde{F}_l(\mathbf{k}) e^{i\mathbf{k} \cdot \mathbf{x}} \quad (1.85)$$

where $\tilde{\mathbf{F}}(\mathbf{k})$ is the spatial Fourier transform of $\mathbf{F}(\mathbf{x})$. This relation allows to deduce explicit expressions for the transverse δ -function.

Note that in the wave equation (1.7) for the vector potential, we have an alternative relation between the current density \mathbf{j} and its transverse part:

$$\mathbf{j}_\perp = \mathbf{j} - \varepsilon_0 \partial_t \nabla \phi,$$

The ‘longitudinal part’ that is subtracted here is thus related to the electrostatic field created by the corresponding charge density. More details are discussed in the exercises.

1.3 Photons and the quantum vacuum

1.3.1 ‘Photons’

The present quantized description of the electromagnetic field allows us to give a more precise meaning to the word ‘photon’. *A photon is an ex-*

citation of a mode of the field. We have seen that the quantized field reduces to a collection of harmonic oscillators, one for each mode. As we know from the harmonic oscillator, its stationary states are labelled by non-negative numbers $n = 0, 1, \dots$. One says that in these states, the mode contains ‘ n photons’. The creation operator a^\dagger whose action on these states is $a^\dagger|n\rangle \propto |n+1\rangle$, thus ‘creates one photon’. This picture is consistent with the assumption (dating back to Einstein (1905)) that photons correspond to ‘energy packets’ of $\hbar\omega_k$ of the electromagnetic field. If a plane wave mode expansion is used, we can also say that the momentum of a photon is $\hbar\mathbf{k}$, as we know from de Broglie (1926) or from the Compton effect.

It is however possible to use different mode expansions for the same field. For example, we could have used an expansion in terms of spherical vector harmonics which differs from the plane wave expansion by a (infinite-dimensional) unitary transformation. A single-photon state in the plane wave basis thus becomes a superposition of single-photon states in infinitely many spherical modes. Conversely, a ‘photon’ in this description would not correspond to a plane wave (its momentum would not be $\hbar\mathbf{k}$), but it would have a definite angular momentum with respect to the origin. It is even possible to define photons that are wavepackets localized in time, by superposing plane waves with neighboring frequencies. This picture allows to describe experiments with ‘single-photon pulses’. We refer to the exercises to look at these properties in more detail.

Finally, we precall that the computation of a suitable set of mode functions is a ‘classical’ problem: no quantum theory is needed to state it. Using a cubic ‘quantization’ box, one can show that the plane waves with wave vectors $\mathbf{k} = (2\pi/L)(n_x, n_y, n_z)^T$ ($n_i \in \mathbb{Z}$) are orthogonal with respect to the scalar product

$$\int_{L^3} d^3x \mathbf{F}^*(\mathbf{x}) \cdot \mathbf{G}(\mathbf{x}). \quad (1.86)$$

This is clear for different wave vectors, $\mathbf{k} \neq \mathbf{k}'$. But for a given \mathbf{k} , one can also find two orthogonal polarization vectors $\boldsymbol{\varepsilon}_{1,2}$ that give orthogonal modes (the scalar product $\boldsymbol{\varepsilon}_1^* \cdot \boldsymbol{\varepsilon}_2$ is zero). The corresponding magnetic field mode functions are also orthogonal. The electrodynamics becomes a ‘quantum’ theory only when the amplitudes of the mode functions become suitably normalized operators.

1.3.2 Vacuum fluctuations

In the vacuum state, the expectation value $\langle \mathbf{E}(\mathbf{x}, t) \rangle = 0$ because a_κ annihilates the vacuum state and a_κ^\dagger can be made to act to the left on the vacuum state which is annihilated.

Note. The same is true for any stationary (or number) state, see Sec.2.2 below.

The vacuum fluctuations become visible in the next moment of the field: for a given mode κ ,

$$\begin{aligned} \frac{\hbar\omega_\kappa}{2\varepsilon_0} \langle (\mathbf{f}_\kappa(\mathbf{x})a(t) + \text{h.c.})^2 \rangle &= \frac{\hbar\omega_\kappa}{2\varepsilon_0} |\mathbf{f}_\kappa(\mathbf{x})|^2 \langle a_\kappa(t)a_\kappa^\dagger(t) \rangle \\ &= \frac{\hbar\omega_\kappa}{2\varepsilon_0} |\mathbf{f}_\kappa(\mathbf{x})|^2 \end{aligned} \quad (1.87)$$

in the last step, we have used that $a_\kappa(t) = a_\kappa(0) e^{-i\omega_\kappa t}$. In the case of plane wave modes, we have $|\mathbf{f}_\kappa(\mathbf{x})|^2 = 1/V$. The sum over wave vectors \mathbf{k} and polarization indices λ can be written in the form

$$\langle \mathbf{E}^2(\mathbf{x}, t) \rangle = \int_0^\infty \frac{d\omega}{2\pi} \frac{\hbar\omega_\kappa}{2\varepsilon_0} \rho(\omega) \quad (1.88)$$

where $\rho(\omega)$ is the so-called ‘density of modes’ (per unit frequency and unit volume):

$$\rho(\omega) = \frac{2\pi}{V} \sum_{\mathbf{k}\lambda} \delta(\omega - \omega_{\mathbf{k}\lambda}) \quad (1.89)$$

For the plane wave modes, $\omega_{\mathbf{k}\lambda}$ only depends on the magnitude of \mathbf{k} , and in the continuum limit,

$$\sum_{\mathbf{k}} = V \int \frac{d^3k}{(2\pi)^3} \quad (1.90)$$

we get after a simple integration

$$\rho(\omega) = \frac{2}{\pi} \frac{\omega^2}{c^3} \quad (1.91)$$

The full vacuum fluctuation of the electric field is then infinite

$$\langle \mathbf{E}^2(\mathbf{x}, t) \rangle = \frac{\hbar}{\pi\varepsilon_0 c^3} \int_0^\infty \frac{d\omega}{2\pi} \omega^3 \quad (1.92)$$

because the integral diverges at the upper limit. This correlates with an infinite electromagnetic energy density in vacuum (multiply with $\varepsilon_0/2$ and add the magnetic component, which doubles the result). This “infinite vacuum energy” is one of the unresolved problems in physics. “Intuitive cutoffs” at short wavelengths, for example, at the Planck scale (10^{-32} m) give a finite energy density, but with a value that differs by something like 100 orders of magnitude from the energy density associated with cosmological observations (including “dark energy”, “cosmological constants” and so on).

Exercise. Find a cutoff such that the vacuum energy density equals the ‘critical density of the Universe’ (the critical mass density is roughly 10^{-29} g/cm³, given the current expansion rate of the Universe.)

A finite value can be found, if we compute an autocorrelation function of the electric field. A similar calculation gives

$$\langle \mathbf{E}(\mathbf{x}, t) \cdot \mathbf{E}(\mathbf{x}, t') \rangle = \int_0^\infty \frac{d\omega}{2\pi} \frac{\hbar\omega_\kappa}{2\varepsilon_0 c^3} \rho(\omega) e^{-i\omega(t-t')} \quad (1.93)$$

This leads to the integral representation of the Γ -function in the complex plane and finally to

$$\langle \mathbf{E}(\mathbf{x}, t) \cdot \mathbf{E}(\mathbf{x}, t') \rangle = \frac{3\hbar}{\pi^2 \varepsilon_0 c^3 \tau^4} \quad (1.94)$$

which is finite for all $\tau = t - t' \neq 0$. For any finite value of τ , the electric vacuum energy is thus of the order of $\hbar\omega_\tau/\lambda_\tau$ with the characteristic frequency $\omega_\tau = 1/\tau$ and wavelength $\lambda_\tau = c\tau$.

1.3.3 Casimir energy

The Casimir force is the attraction between two metallic mirrors placed in vacuum. It is interpreted in terms of the change in the zero-point energy (the famous $\frac{1}{2}\hbar\omega$ of the harmonic oscillator ground state) induced by the presence of the mirrors. We give here an sketch of the calculation done by Casimir around 1948 [Proc. Kon. Ned. Akad. Wet. 51 (1948) 793].

We consider the ground state energy of the multi-mode electromagnetic field

$$E_0 = \sum_{\mathbf{k}\lambda} \frac{\hbar\omega_{\mathbf{k}\lambda}}{2}$$

that is of course infinite and compare the cases of a planar cavity formed by two mirrors (distance L) and empty space (i.e., two mirrors infinitely apart). In the first case, we have standing wave modes between the mirrors with a frequency

$$\omega_n = c\sqrt{K^2 + k_n^2}, \quad k_n = \frac{n\pi}{L} \quad (1.95)$$

with $K^2 = k_x^2 + k_y^2$ and $n = 1, 2, \dots$, while in empty space,

$$\omega = c\sqrt{K^2 + k_z^2}$$

with $-\infty < k_z < \infty$. We first compute the difference in the electromagnetic mode density per volume AL where A is the ‘quantization area’ in the xy -plane. We cheat with the polarizations and multiply by a factor 2:

$$\begin{aligned} \rho_L(\omega) &= \frac{4\pi}{AL} \sum_{\mathbf{K}, n} \delta\left(\omega - c\sqrt{K^2 + k_n^2}\right) \\ &= \frac{2}{L} \int_0^\infty K dK \sum_n \delta\left(\omega - c\sqrt{K^2 + k_n^2}\right) \end{aligned} \quad (1.96)$$

The integration over K can be performed with the substitution $K \mapsto c\sqrt{K^2 + k_n^2}$ and gives

$$\rho_L(\omega) = \frac{2\omega}{Lc^2} \sum_n \Theta(\omega - ck_n) \quad (1.97)$$

where Θ is the step function. It arises because for a given n , there are no modes with frequency smaller than ck_n , see Eq.(1.95). The same calculation in the infinite volume gives

$$\rho_\infty(\omega) = \frac{2\omega}{c^2} \int_0^\infty \frac{dk_z}{\pi} \Theta(\omega - ck_z) \quad (1.98)$$

The k_z integral can of course be performed, but we keep it here to illustrate one of the basic features of the Casimir calculation: the result originates from the difference between a ‘discrete spectrum’ (the sum over the k_n) and a continuum (the integral over k_z).

The Casimir energy is now found as the difference in vacuum energy *per area* in the space of length L between the mirrors:

$$\Delta E = L \int_0^\Omega \frac{\hbar\omega}{2} (\rho_L(\omega) - \rho_\infty(\omega)) \quad (1.99)$$

We have introduced an upper cutoff frequency Ω because the integrals are likely to diverge in the UV. One of the mathematical difficulties (that we are not going to discuss here) is to what extent the results depend on the cutoff. At a suitable stage of the calculation, we are going to take the limit $\Omega \rightarrow \infty$, of course.

The ω -integrals can be performed before the sum over n (the integral over k_z), and one gets

$$\Delta E = \frac{\hbar}{6\pi c^2} \left[\sum_{n=1}^{\lfloor \Omega L/\pi c \rfloor} (\Omega^3 - (ck_n)^3) - L \int_0^{\Omega/c} \frac{dk_z}{\pi} (\Omega^3 - (ck_z)^3) \right] \quad (1.100)$$

where $\lfloor x \rfloor$ is the largest integer smaller than x . Introducing the number $N = \Omega L/\pi c$ and the dimensionless integration variable $z = k_z L/\pi$, this can be written in the form

$$\Delta E = \frac{\hbar c \pi^2}{6L^3} \left[\sum_{n=1}^{\lfloor N \rfloor} (N^3 - n^3) - \int_0^N dz (N^3 - z^3) \right] \quad (1.101)$$

The difference in brackets is some magic number and equal to $-1/120$ in the limit $N \rightarrow \infty$. (A proof is sketched below). The Casimir energy of two mirrors is thus equal to

$$\Delta E = -\frac{\hbar c \pi^2}{720L^3} \quad (1.102)$$

so that the force per unit area is $F_C/A = -\hbar c \pi^2/240L^4$. Since the energy decreases as $L \rightarrow 0$, the two mirrors placed in vacuum attract each other.

Note that this result is independent of the nature of the mirrors, as well as their electric charge. The electromagnetic field only enters inasmuch as its modes give a contribution to the energy of the vacuum state. Field theorists have computed the contribution to the Casimir energy from the Dirac electron field, for example. It is small if the mirror separation is large compared to the Compton wavelength $\hbar/mc \approx 2.5$ pm — which is nearly always the case. The Casimir energy, being attractive, is sometimes thought of a means to ‘stabilize’ a classical model of the electron (a bag of charge) against the Coulomb repulsion. The sign of the Casimir force depends, however, on the boundary conditions one takes at the mirror surfaces. ‘Perfectly permeable’ mirrors (that give von Neumann instead of Dirichlet conditions for the tangential electric field) also attract each other,




Figure 1.1: Integration contour for (1.104).

but a ‘mixed pair’ (perfect Dirichlet facing von Neumann) gives repulsion. This is discussed in a pedagogical way in a paper by Hushwater, “Repulsive Casimir force as a result of vacuum radiation pressure” [Am. J. Phys. 65 (1997) 381].

Sum minus integral

We use a trick in the complex plane. There is a theorem for functions f and D that are analytical in a domain limited by the integration contour \mathcal{C} :

$$\frac{1}{2\pi i} \oint_{\mathcal{C}} dz f(z) \frac{d}{dz} \log D(z) = \sum_n f(z_n) \quad (1.103)$$

where the z_n are the zeros of D in the interior of the contour. We will use $f(z) = N^3 - z^3$ and choose $D(z)$ such that it is zero for the values $z_n = n$: $D(z) = \sin(\pi z)$. The differentiation under the integral sign gives

$$\sum_{n=0}^N (N^3 - n^3) = \frac{1}{2} \oint_{\mathcal{C}_N} dz (N^3 - z^3) \frac{e^{i\pi z} + e^{-i\pi z}}{e^{i\pi z} - e^{-i\pi z}} \quad (1.104)$$

We chose an integration contour as shown in fig. 1.1 running from $+N$ above the real axis to 0 and going back to $+N$ below the real axis (the sum over all positive zeros of $\sin \pi z$ thus gives the sum on the left hand side). We are eventually interested in the limit $N \rightarrow \infty$. Make the following

transformations on the upper and lower part of the contour:

$$\begin{aligned} \text{upper part: } & \frac{e^{i\pi z} + e^{-i\pi z}}{e^{i\pi z} - e^{-i\pi z}} = -1 + \frac{2e^{i\pi z}}{e^{i\pi z} - e^{-i\pi z}} \\ \text{lower part: } & \frac{e^{i\pi z} + e^{-i\pi z}}{e^{i\pi z} - e^{-i\pi z}} = 1 + \frac{2e^{-i\pi z}}{e^{i\pi z} - e^{-i\pi z}} \end{aligned}$$

The constants ± 1 give for both the upper and lower path an integral over $N^3 - z^3$ that can be combined into

$$\sum_{n=0}^N (N^3 - n^3) = \int_0^N dz (N^3 - z^3) + \oint_{\mathcal{C}} dz (N^3 - z^3) \frac{e^{\pm i\pi z}}{e^{i\pi z} - e^{-i\pi z}}$$

In the second integral, the exponential takes the appropriate sign on the upper and lower parts of the contour. The first integral on the right hand side is exactly the integral that we have to subtract in Eq.(1.101). The upper and lower parts of the contour can now be shifted onto the (positive or negative) imaginary axis because the integrand has no singularities (these are located on the real axis only). The quarter-circles with radius $|z| = N$ contribute exponentially small amounts for large N because of the $e^{\pm i\pi z}$.

Choosing $z = \pm it$ on the imaginary axis, the integrals there give

$$\begin{aligned} & \oint_{\mathcal{C}} dz (N^3 - z^3) \frac{e^{\pm i\pi z}}{e^{i\pi z} - e^{-i\pi z}} \\ &= i \int_0^\infty dt \frac{[N^3 - (it)^3] e^{-\pi t}}{e^{\pi t} - e^{-\pi t}} - i \int_0^\infty dt \frac{[N^3 - (-it)^3] e^{-\pi t}}{e^{\pi t} - e^{-\pi t}} \\ &= -2 \int_0^\infty dt \frac{t^3}{e^{2\pi t} - 1} \end{aligned}$$

Note that the imaginary parts of the two integrals that involve N^3 cancel each other: we have finally eliminated the cutoff.

You may have encountered the last integral in the context of blackbody radiation. Changing to the integration variable $t' = 2\pi t$, the integral gives $1/240$, so that we have in the end

$$\lim_{N \rightarrow \infty} \left(\sum_{n=0}^N (N^3 - n^3) - \int_0^N dz (N^3 - z^3) \right) = -\frac{2}{240} = -\frac{1}{120} \quad (1.105)$$

as announced in the text.

Chapter 2

Quantum states of the radiation field

2.1 Observables

The quantized light field can be in different states. We start here with a single mode of the field. This may be a oversimplification, but single-mode fields have become part of the experimental reality with the advent of high-quality optical cavities. These devices give an electromagnetic field whose amplitude, in the region between two well-reflecting mirrors, is much higher at some resonant frequencies. The ‘mode function’ is in this case not a plane wave, of course, but a standing wave. In the transverse directions, one often has a gaussian profile. Around a cavity resonance, it is a frequent approximation to treat the full field as if it contained only a single mode. The coupling to other modes may be taken into account as a loss.

The electric field is given by

$$\mathbf{E}(\mathbf{x}, t) = E_1 \boldsymbol{\varepsilon} \left(a(t) + a^\dagger(t) \right) \sin kz \quad (2.1)$$

where z is the coordinate along the cavity axis and $k = n_z \pi / L$. The factor E_1 can be called the ‘electric field per photon’. From the multimode expansion, E_1 is given by the prefactor $E_k = (\hbar \omega_k / 2 \varepsilon_0 V)^{1/2}$. The corresponding ‘intensity’ is

$$I_{1\text{ph}} = \varepsilon_0 c E_k^2 = \frac{\hbar \omega_k c}{2V}. \quad (2.2)$$

In a cavity, we can take for V the volume ‘filled’ by the mode. For a transverse mode size of 1 micrometer and a cavity length of 1 cm, we get $I_{1\text{ph}} \sim 10^3 \text{ mW/cm}^2$ which is not really small. The total power, however, is quite small: about 10^{-8} W . Note also that these numbers are based on very ‘tight’ (diffraction-limited) focussing — beams with larger cross-section have a smaller ‘field per photon’.

In the Heisenberg picture, the field operator evolves as

$$\mathbf{E}(\mathbf{x}, t) = E_1 \boldsymbol{\epsilon} \left(a e^{-i\omega t} + a^\dagger e^{-i\omega t} \right) \sin kz \quad (2.3)$$

A combination of annihilation and creation operators like the one in parentheses is called a ‘quadrature’. Quadratures always come in pairs. One can find a second quadrature variable by shifting the origin of time by one quarter period: $\propto -i a e^{-i\omega t} + i a^\dagger e^{i\omega t}$. This corresponds to the magnetic field (compare eqs. 1.64 and 1.65). In analogy to the harmonic oscillator, one often uses the following quadrature variables

$$X = \frac{a + a^\dagger}{\sqrt{2}} \quad P = \frac{a - a^\dagger}{\sqrt{2}i} \quad (2.4)$$

The simplest quantum states of the single mode field are given by the well-known stationary states of the harmonic oscillator.

2.2 Number states

These quantum states are called ‘Fock states’ or ‘number states’ $|n\rangle$. They are eigenstates of the ‘photon number operator’

$$\hat{n} = a^\dagger a = aa^\dagger - 1 \quad (2.5)$$

and are generated by applying the creation operator to the ground state of the oscillator:

$$|n\rangle = \frac{1}{\sqrt{n!}} (a^\dagger)^n |0\rangle \quad (2.6)$$

The expectation value of the annihilation operator is zero in a number state:

$$\langle a \rangle_n = \langle n | a | n \rangle = \sqrt{n} \langle n | n - 1 \rangle = 0 \quad (2.7)$$

The same is true for the creation operator. It follows that the electric field average vanishes in a Fock state:

$$\langle \mathbf{E}(\mathbf{x}, t) \rangle_n = 0 \quad (2.8)$$

The ground state of the field mode is called the ‘vacuum’ (no photon, i.e., no excitation present). It is found by looking for the state that is annihilated by the annihilation operator: $a|\text{vac}\rangle = 0$. Obviously, this is also an eigenstate of the photon number operator with zero photons: $|\text{vac}\rangle = |0\rangle$. In the vacuum state, the electric field is also zero on average, of course.

But there are fluctuations around this average, called ‘quantum noise’. In the vacuum state of the single mode (2.1), e.g., we get

$$\langle \mathbf{E}(\mathbf{x}, t)^2 \rangle_0 = E_1^2 \sin^2 kz \langle 0 | (a(t) + a^\dagger(t)) (a(t) + a^\dagger(t)) | 0 \rangle \quad (2.9)$$

and this combination of operators gives an average

$$\langle 0 | (a(t) + a^\dagger(t)) (a(t) + a^\dagger(t)) | 0 \rangle = \langle 0 | a(t) a^\dagger(t) | 0 \rangle = 1 \quad (2.10)$$

The ‘vacuum noise’ in our mode is thus given by the squared single photon field $E_1^2 \sin^2 kz$. Similarly, the other quadrature variable $a(t) - a^\dagger(t)$ shows a noise strength of unity. This is in accordance with Heisenberg’s indeterminacy relation, since

$$[a(t) + a^\dagger(t), a(t) - a^\dagger(t)] = -2. \quad (2.11)$$

In the exercises, you are asked to compute the variances of these quadratures in an arbitrary number state $|n\rangle$.

The electric field fluctuations in the vacuum state play an important role in the interpretation of the Casimir effect and of the spontaneous decay of excited atomic states.

Finally, the creation and annihilation operators connect states whose photon numbers differ by one. In this sense, the ‘creation operator’ a^\dagger creates one photon since for example

$$\langle 1 | a^\dagger | 0 \rangle = 1. \quad (2.12)$$

This matrix element plays an important role when one computed the probability amplitude that an excited atomic state emits a photon. For stimulated emission, one needs $\langle n+1 | a^\dagger | n \rangle = \sqrt{n+1}$. Similarly, the ‘annihilator’ a destroys one photon:

$$\langle 0 | a | 1 \rangle = 1. \quad (2.13)$$

This matrix element is needed to compute absorption, and in the general case, $\langle n-1|a|n\rangle = \sqrt{n}$.

2.3 Thermal states

This class of field states is also quite important, but it is more general than the ‘pure’ states described so far. In a thermal state, one has to use both classical and quantum statistics. The easiest way to deal with that is to use stationary states, hence the number states we just found. In this case, one can use Boltzmann statistics to describe the field at thermal equilibrium.

Each stationary state $|n\rangle$ can occur with a classical probability proportional to the Boltzmann factor $e^{-n\hbar\omega/k_B T}$. The normalization of these probabilities (the ‘partition function’) is a simple geometrical series

$$Z = \sum_{n=0}^{\infty} e^{-n\hbar\omega/k_B T} = \frac{1}{1 - e^{-\hbar\omega/k_B T}}, \quad (2.14)$$

so that the probability to find n photons in a single mode is given by

$$p_n(T) = (1 - e^{-\hbar\omega/k_B T}) e^{-n\hbar\omega/k_B T} \quad (2.15)$$

At optical frequencies and room temperature, the exponential has a large negative argument for $n \geq 1$ so that the field is essentially at zero temperature. This is different for microwave radiation, e.g., or for star atmospheres.

Simple exercise: mean photon number. Result:

$$\langle \hat{n} \rangle_T = \frac{1}{e^{\hbar\omega/k_B T} - 1}. \quad (2.16)$$

Exercise: photon number variance. Result:

$$(\Delta n)_T^2 = \frac{e^{\hbar\omega/k_B T}}{(e^{\hbar\omega/k_B T} - 1)^2} = \frac{1}{4 \sinh^2(\hbar\omega/2k_B T)}. \quad (2.17)$$

Electric field fluctuations in a single mode at finite temperature:

$$\langle \mathbf{E}^2(\mathbf{x}, t) \rangle_T = E_1^2 \sin^2 kz \langle a(t)a^\dagger(t) + a^\dagger(t)a(t) \rangle_T = E_1^2 \sin^2 kz (2\langle \hat{n} \rangle_T + 1) \quad (2.18)$$

they are enhanced by a factor $2\langle \hat{n} \rangle_T + 1 = \coth(\hbar\omega/2k_B T)$ compared to zero temperature.

Finally, the thermal state is often expressed in terms of a *density operator* $\hat{\rho}$. This is an operator on the space of number states. In thermal equilibrium, its expansion in the number state basis can be read off from the probabilities (2.15)

$$\hat{\rho} = \sum_{n=0}^{\infty} p_n(T) |n\rangle\langle n| \quad (2.19)$$

Average values with respect to both quantum and classical statistics can now be obtained from a trace of the density operator. For the photon number operator, e.g.,

$$\langle \hat{n} \rangle_{\rho} = \sum_{n=0}^{\infty} n p_n(T) = \sum_{n=0}^{\infty} \langle n | \hat{n} \hat{\rho} | n \rangle = \text{tr}(\hat{n} \hat{\rho}). \quad (2.20)$$

For an arbitrary operator A , this is generalized to

$$\langle \hat{A} \rangle_{\rho} = \text{tr}(\hat{A} \hat{\rho}) = \sum_{n=0}^{\infty} \langle n | \hat{A} \hat{\rho} | n \rangle \quad (2.21)$$

The advantage of this formalism is that the trace can be taken in any basis, and one can choose a basis adapted to the operator whose average one is interested in. The second advantage of using the density matrix is that it gives a suitable description of a quantum system whose dynamics is not completely known and can only be specified by probabilities. In that case, one formulates an equation of motion for the density matrix from the solution of which the averages of all interesting quantities can be calculated.

2.4 Coherent states

2.4.1 Basic properties

The coherent state $|\alpha\rangle$ is an eigenstate of the annihilation operator:

$$a|\alpha\rangle = \alpha|\alpha\rangle \quad (2.22)$$

Since a is not an hermitean operator, α can be complex. *In a coherent state, the average electric field is nonzero:*

$$\langle \mathbf{E}(\mathbf{x}, t) \rangle_{\alpha} = E_1 \sin kz \langle \alpha | (a(t) + a^{\dagger}(t)) | \alpha \rangle = E_1 \sin kz (\alpha e^{-i\omega t} + \alpha^* e^{i\omega t}). \quad (2.23)$$

We have assumed the field in a coherent state of the initial annihilator a . This expression is the same that we have used in chapter 1 for a classical, monochromatic field. The magnetic field quadrature also has on average its classical value in a coherent state. Coherent states are thus very useful to represent laser fields. We see that α measures the electric field strength in units of the ‘single photon field’ E_1 . If we compute the average photon number in a coherent state, we get

$$\langle \hat{n} \rangle_\alpha = \langle a | a^\dagger a | \alpha \rangle = |\alpha|^2, \quad (2.24)$$

so that as an order of magnitude $\langle E \rangle \approx E_1 \langle \hat{n} \rangle^{1/2}$ (note the nonlinear dependence).

Coherent states are not stationary, but rotate in the complex α -plane: if $|\psi(0)\rangle = |\alpha\rangle$, then $|\psi(t)\rangle = |\alpha e^{-i\omega t}\rangle$. This can be shown using the expansion of a coherent state in terms of number states:

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle \quad (2.25)$$

Note that number states with arbitrarily high photon numbers are present in a coherent state. More specifically, we can introduce the probability $p_n(\alpha)$ of finding n photons in a coherent state:

$$p_n(\alpha) = |\langle n | \alpha \rangle|^2 = e^{-|\alpha|^2} \frac{|\alpha|^{2n}}{n!} \quad (2.26)$$

which is a ‘Poisson distribution’ (the probability distribution of the sum of independent random bits). The average photon number

$$\begin{aligned} \langle \hat{n} \rangle &= \langle \alpha | a^\dagger a | \alpha \rangle \\ &= \alpha^* \alpha \langle \alpha | \alpha \rangle = |\alpha|^2 \end{aligned} \quad (2.27)$$

and its fluctuations are found from

$$\begin{aligned} \langle \hat{n}^2 \rangle &= \langle \alpha | a^\dagger a a^\dagger a | \alpha \rangle \\ &= \langle \alpha | (a^\dagger a^\dagger a a + a^\dagger a) | \alpha \rangle \\ &= |\alpha|^4 + |\alpha|^2 \end{aligned} \quad (2.28)$$

Hence, the relative fluctuation is small as the coherent state amplitude increases: $(\Delta n)^2 / \langle \hat{n} \rangle^2 = 1/|\alpha|^2$.




Figure 2.1: Representation of a coherent state in phase space.

2.4.2 Phase space representation

The field quadratures also show quantum fluctuations around their classical average in a coherent state. This is inevitable because of the Heisenberg inequality. In the exercises, you are asked to show that these are equal to the quantum noise in the vacuum state (which is in fact a particular coherent state with $\alpha = 0$). This result is often graphically displayed in the complex α -plane by the sketch shown in fig. 2.1. Note that since $a = (X + iP)/\sqrt{2}$, the α -plane may be identified with the classical phase space of a harmonic oscillator. The gray area in this sketch indicates values for the position and momentum quadratures that are probable outcomes of measurements. This representation is of course schematic since X and P cannot be measured simultaneously. We can give it a precise meaning by considering the following function

$$Q_\rho(\alpha) = \langle \alpha | \hat{\rho} | \alpha \rangle \quad (2.29)$$

This is called the ‘Husimi-’ or ‘Q-function’ and is known as a quasi-probability distribution on the phase space. The Q-function is defined on the entire α -plane, and it is positive there. Loosely speaking, we can interpret it as the probability to find the coherent state $|\alpha\rangle$ when the field mode is measured. (There are some subtleties related to the fact that coherent states are not eigenstates of an hermitean operator, hence this is not directly measurable. Several other distribution functions can also be defined.)

For a coherent state $|\beta\rangle$, the Q-function is

$$Q_\beta(\alpha) = \exp(-|\alpha - \beta|^2)$$

hence an isotropic gaussian centered around $\alpha = \beta$ with a width of order unity. The contour lines of this function are sketched in Fig.2.1. The finite width around the maximum takes into account the ‘quantum fluctuations’ of the state.

How would a number state look in this sketch? A first guess is a ‘ring’, since the photon number (or energy) is fixed and shows no fluctuations. This is indeed true, because

$$Q_n(\alpha) = |\langle n|\alpha\rangle|^2 = \frac{e^{-|\alpha|^2} |\alpha|^{2n}}{n!}$$

if we consider this as a function of α , it increases from zero like a power law $|\alpha|^{2n}$ and is cut off for large $|\alpha|$ by the exponential. The maximum is around $|\alpha| = \sqrt{n}$ where number state $|n\rangle$ and coherent state $|\alpha\rangle$ have the same (average) photon number. Of course, the distribution function does not depend on the angle in the α -plane.

Exercise. Q-function of a thermal state.

2.4.3 Classical sources generate coherent states

Finally, how is it possible to generate a coherent state physically? To this end, consider the single-mode Hamiltonian

$$H = \hbar\omega \left(a^\dagger a + \frac{1}{2} \right) + \hbar \left(g e^{-i\omega_L t} + g^* e^{i\omega_L t} \right) \left(a + a^\dagger \right) \quad (2.30)$$

The interaction describes, e.g., the coupling of a classical oscillating current source $j(t)$ with the field mode. One can check that this Hamiltonian generates the same time evolution as the full Maxwell equation by starting from the wave equation for the quantized vector potential. The current density $\mathbf{j}_\perp(\mathbf{x}, t)$ needs to be projected onto the mode function $\mathbf{f}_\kappa(\mathbf{x})$ of the single mode a in question; this gives the term $g \left(e^{-i\omega_L t} + e^{i\omega_L t} \right)$ for an oscillating current at the frequency ω_L .

2.4.4 Resonance approximation

To solve the time-dependent Schrödinger equation, we transform in the so-called interaction picture:

$$|\psi(t)\rangle = e^{-iH_0t/\hbar}|\tilde{\psi}(t)\rangle$$

where H_0 describes the ‘free evolution’ at ω . We find that $|\tilde{\psi}(t)\rangle$ solves a Schrödinger equation with a Hamiltonian

$$H_{\text{int}}(t) = e^{iH_0t/\hbar} H e^{-iH_0t/\hbar} - H_0 \quad (2.31)$$

$$= \hbar \left(g e^{-i\omega_L t} + g^* e^{i\omega_L t} \right) \left(a e^{-i\omega t} + a^\dagger e^{i\omega t} \right) \quad (2.32)$$

This Hamiltonian is still time-dependent. We now make the ‘resonance approximation’, assuming that the source frequency ω_L is near the mode frequency ω . In that case, we identify two classes of terms in Eq.(2.32):

- ‘resonant terms’ that vary slowly on a time scale $1/\omega$:

$$g^* a e^{-i(\omega-\omega_L)t} + g a^\dagger e^{i(\omega-\omega_L)t}$$

- ‘non-resonant terms’ that show rapid oscillations at a frequency near 2ω :

$$g a e^{-i(\omega_L+\omega)t} + g^* a^\dagger e^{i(\omega_L+\omega)t}$$

This has crucial consequences for the time-evolution operator that corresponds to $H_{\text{int}}(t)$: it involves the integral of $H_{\text{int}}(t)$ with respect to time. On time scales larger than $1/\omega$, the non-resonant terms are averaged away, and only the slowly-varying resonant terms give a contribution.

This motivates the so-called *resonance approximation* for the Hamiltonian (2.33): we replace it by

$$H = \hbar\omega \left(a^\dagger a + \frac{1}{2} \right) + \hbar \left(g a^\dagger e^{-i\omega_L t} + g^* a e^{i\omega_L t} \right) \quad (2.33)$$

throwing away those terms that give non-resonant terms in the interaction picture. This approximation also has a clear interpretation when the source is quantized. In that case, $g e^{-i\omega_L t}$ is replaced by an annihilation operator b , and the time evolution of b is generated by a harmonic oscillator Hamiltonian at frequency ω_L . Now, the resonant terms in the interaction Hamiltonian correspond to

- photon (excitation quantum) transfer from source to mode: ba^\dagger , or the other way round: $b^\dagger a$;
- the non-resonant terms correspond to the simultaneous creation of two excitations, one in the source, one in the mode: $b^\dagger a^\dagger$, or simultaneous disappearance of two excitations: ba . By a (naive) application of energy conservation, these processes are ‘forbidden’.

Note that there is actually no contradiction: at sufficiently short interaction times, the time-energy ‘uncertainty relation’ permits the non-resonant processes to ‘happen virtually’. Energy conservation in the formulation of photon number conservation only applies on sufficiently long time scales (larger than the period $2\pi/\omega$).

The resonance approximation is often called in quantum optics the *rotating wave approximation* for historical reasons.

2.4.5 The resonant case

We now focus on exact resonance between the source and the mode, $\omega_L = \omega$. In that case, the interaction representation leads to a Hamiltonian in the resonance approximation that is no longer time-dependent. The time evolution operator is thus easy to find out and of the form

$$U(t) = D(\alpha) \equiv \exp(\alpha a^\dagger - \alpha^* a)$$

with $\alpha = -igt$. Starting with the mode in the vacuum state, we thus find using the Baker-Hausdorff-Campbell identity (exercise!)

$$|\psi(t)\rangle = \exp(\alpha a^\dagger - \alpha^* a) |0\rangle = e^{-|\alpha|^2/2} e^{\alpha a^\dagger} e^{-\alpha^* a} |0\rangle \quad (2.34)$$

Now the annihilation operator gives 0 when acting on the vacuum state, so that its exponential reduces to unity here. Expanding the exponential with the creation operator in a power series, we find

$$|\psi(t)\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{(\alpha)^n}{\sqrt{n!}} |n\rangle = |\alpha\rangle \quad (2.35)$$

This interaction thus generates a coherent state with amplitude $\alpha = -igt$ that grows linearly in time. To obtain a stationary result, either the ‘oscillator amplitude’ g can be made time-dependent (detuning slightly from resonance), or loss processes have to be added.

2.4.6 Displacement operator

We have just shown that coherent states can be obtained by applying a ‘displacement operator’ to the vacuum state:

$$|\alpha\rangle = D(\alpha)|0\rangle \quad D(\alpha) = \exp\{\alpha a^\dagger - \alpha^* a\} \quad (2.36)$$

This unitary operator also displaces the creation and annihilation operators as follows (prove it yourself)

$$D^\dagger(\alpha) a D(\alpha) = a + \alpha \quad (2.37)$$

$$D^\dagger(\alpha) a^\dagger D(\alpha) = a^\dagger + \alpha^*. \quad (2.38)$$

This identity is useful to show that the field quadrature fluctuations in a coherent state are those of the vacuum state.

2.5 Squeezed states

You should have got the feeling up to now that the quantized field essentially differs from a classical field by its (‘quantum’) fluctuations. So people have thought whether it is possible to reduce the quantum noise in a field quadrature to get something even ‘more classical’ – or having less noise. This can be achieved in part, to 50%, say. Of course, one cannot beat the Heisenberg inequality, and the reduced fluctuations in one quadrature have to be paid by enhanced noise in the other one.

In the exercises, you are asked to show that the following operator

$$S(\xi) = \exp(\xi a^{\dagger 2} - \xi^* a^2) \quad (2.39)$$

‘squeezes’ the fluctuations of the vacuum state such that one quadrature component (fixed by the phase of the complex number ξ) has less quantum noise. Note that a ‘general quadrature’ variable may be defined by

$$X_\theta = \frac{a e^{i\theta} + a^\dagger e^{-i\theta}}{\sqrt{2}} \quad (2.40)$$

Two ‘orthogonal’ quadratures are then given by X_θ and $X_{\theta+\pi/2}$. The usual position and momentum quadratures correspond to $\theta = 0, \pi/2$. In the state $S(\xi)|0\rangle$ with $\xi = |\xi| e^{-2i\theta}$, the uncertainties of the quadratures are

$$\Delta X_\theta^2 = \frac{e^{2|\xi|}}{2}, \quad \Delta X_{\theta+\pi/2}^2 = \frac{e^{-2|\xi|}}{2}, \quad (2.41)$$

their product being unchanged.

Note also that the ‘squeezed vacuum’ $S(\xi)|0\rangle$, when expanded in the Fock basis, contains only even photon number states:

$$S(\xi)|0\rangle = \sum_{m=0}^{\infty} c_{2m}|2m\rangle. \quad (2.42)$$

This is because the squeezing operator (2.39) only contains even powers of the creation operator.

Exercise: compute the coefficients c_{2m} . Answer (details may be wrong):

$$c_{2m} = N \left(\frac{(2m-1)!!}{(2m)!!} \right)^{1/2} e^{-2im\phi} \tanh^m r$$

where N is a normalization, $\xi = (r/2)e^{2i\phi}$, and $n!!$ is the product $n(n-2)\cdots$ of all numbers with the same parity up to n .

2.5.1 Squeezing from a nonlinear medium

The ‘squeezing’ operator (2.39) can be realized with the interaction Hamiltonian

$$H_{\text{int}} = i\hbar \left(g e^{-2i\omega t} a^{\dagger 2} - g^* e^{2i\omega t} a^2 \right) \quad (2.43)$$

with the squeezing parameter given by $\xi = \int dt g(t)$. This interaction occurs in nonlinear optics. To get a qualitative understanding, imagine a medium with a field-dependent dielectric constant (‘ $\chi^{(2)}$ nonlinearity’). This is usually forbidden for symmetry reasons, but it happens in some special cases. In the electromagnetic energy density, one has

$$u = \frac{\varepsilon(|\mathbf{E}|)}{2} \mathbf{E}^2 + \frac{1}{2\mu_0} \mathbf{B}^2 \quad (2.44)$$

where the linearization

$$\varepsilon(|\mathbf{E}|) = \varepsilon_0 (1 + n_2 |\mathbf{E}|^2) \approx \varepsilon_0 (1 + 2n_2 |\mathbf{E}|)$$

is often appropriate. In the quantum picture, this gives a contribution to the Hamiltonian with a term of third order in the field:

$$H_3 = \varepsilon_0 n_2 \int_V d^3x |\mathbf{E}(\mathbf{x}, t)|^3 \quad (2.45)$$

Let us now pick out two spatial modes of the field and put one of it into a coherent state $|\alpha e^{-i\omega_p t}\rangle$ with a ‘large’ amplitude $|\alpha| \gg 1$. The index ‘p’ is for ‘pump field’. Let us call the other mode (the ‘quantum’ one) the ‘signal’. The electric field is then

$$\mathbf{E}(\mathbf{x}, t) = E_p a_p \boldsymbol{\varepsilon}_p e^{-i(\omega_p t - \mathbf{k}_p \cdot \mathbf{x})} + E_1 \boldsymbol{\varepsilon} a(t) e^{i\mathbf{k} \cdot \mathbf{x}} + \text{h.c.} \quad (2.46)$$

The interaction Hamiltonian thus generates cross terms of the form¹

$$H_{\text{int}} = \dots + \hbar \left(g e^{-i\omega_p t} a_p a^\dagger{}^2 + g^* e^{i\omega_p t} a_p^\dagger a^2 \right) \quad (2.47)$$

$$\hbar g = 3\varepsilon_0 n_2 E_p E_1 \boldsymbol{\varepsilon}_p \cdot \boldsymbol{\varepsilon}^* \int_V d^3x e^{i(\mathbf{k}_p - 2\mathbf{k}) \cdot \mathbf{x}} \quad (2.48)$$

One often ignores the quantum fluctuations of the pump mode and replaces its annihilation operator a_p by the coherent state amplitude α . The interaction (2.47) then looks quite like our model Hamiltonian (2.43).

The nonlinear squeezing parameter $g\alpha$ is nonzero when the pump and signal modes are ‘phase matched’, i.e., $\mathbf{k}_p = 2\mathbf{k}$. For collinear modes, this is achieved by taking $\omega_p = 2\omega$. The spatial integral actually runs only over the region where the nonlinear index n_2 is different from zero. We also see from (2.47) that one ‘pump photon’ with energy $\hbar\omega_p = 2\hbar\omega$ can ‘decay’ into a pair of signal photons. We already anticipated this behaviour in the number state expansion (2.42).

¹We are actually cheating with the polarization vector $\boldsymbol{\varepsilon}$. An accurate description replaces n_2 by a third-rank tensor that produces a scalar out of three vectors.